



IMAGES IN PAEDIATRICS

Aneurysmal ductus arteriosus. Prenatal diagnosis and evolution



Ductus arterioso aneurismático. Diagnóstico prenatal y evolución

María Erroz Ferrer^{a,*}, Patricia Martínez Olorón^{a,c}, Jorge Suárez Alonso^a, Beatriz Ramos Lacuey^b

^a Unidad de Cardiología Infantil, Hospital Universitario de Navarra, Pamplona, Spain

^b Unidad de Neonatología, Hospital Universitario de Navarra, Pamplona, Spain

^c Departamento de Pediatría, Facultad de Ciencias de la Salud, Universidad Pública de Navarra, Pamplona, Spain

Received 7 August 2024; accepted 23 September 2024

Available online 26 February 2025

We present the case of a Saharawi pregnant woman with an unremarkable family and personal history. In the current pregnancy, there was evidence of intrauterine growth restriction (<3rd percentile) and fetal blood flow redistribution. The prenatal ultrasound conducted at 38⁺1 weeks detected an aneurysmal dilatation of the ductus arteriosus (Figs. 1 and 2), so the decision was made to induce labor and admit the infant to the neonatal unit for monitoring due to the potential complications associated with this condition (including spontaneous rupture, embolism, airway erosion, infection, compression of adjacent structures or even death).^{1,2}

The postnatal echocardiography confirmed the presence of a ductus arteriosus aneurysm, and the rest of the cardiovascular evaluation was normal (Appendix A video). During the stay, the infant remained asymptomatic and the aneurysm decreased progressively in size until full closure at day 23 post birth (previous case series have reported favorable outcomes with resolution within 7–35 days). A genetic study was conducted to rule out aortic and vascular diseases, including disorders of connective tissue (such as Marfan or



Figure 1 Prenatal ultrasound. 2D mode. Three-vessel and trachea view. The "V" shape produced by the aorta and pulmonary artery appears in the top left of the image. Visualization of the ductus arteriosus with a saccular dilatation with an aneurysmal appearance and a transverse diameter of 10 mm. Structures: (1) pulmonary artery, (2) aorta, (3) superior vena cava, (4) ductus, (5) trachea.

DOI of original article:
<https://doi.org/10.1016/j.anpedi.2024.503705>

* Corresponding author.
 E-mail address: merrozf@navarra.es (M. Erroz Ferrer).

Ehlers-Danlos syndrome) and variants in the ACTA2 gene, which have been associated with aneurysmal dilatation of the great vessels and other systemic malformations.³ We ought to underscore connective tissue disease on account




Figure 2 Prenatal ultrasound. 2D mode. Sagittal view. Ductal arch with aneurysmal dilatation.

of the increased incidence of complications associated with it.² In this case, the genetic study was negative. The patient remains asymptomatic to date, with no sonographic abnormalities during the follow-up.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.anpede.2025.503705>.

References

1. López A. Aneurisma del ductus arterioso en el feto. Prog Obstet Ginecol. 2010;53:419-21, <http://dx.doi.org/10.1016/j.pog.2010.03.008>. ISSN 0304-5013.
2. Rondón M, Reyna-Villasmil E. Diagnóstico prenatal de aneurisma del ductus arterioso. Avan Biomed. 2017;6:227-30.
3. Ardhanari M, Colin A, Tekin M, Infante JC, Swaminathan S. Aneurysmal Dilatation of Ductus Arteriosus and Pulmonary Artery in Association With ACTA2 Mutation. World J Pediatr Congenit Heart Surg. 2020;11:NP498–500, <http://dx.doi.org/10.1177/2150135120902120>. Epub 2020 May 26. PMID: 32452246.