



IMAGES IN PAEDIATRICS

Gastric *Helicobacter heilmannii* infection**Infección gástrica por *Helicobacter heilmannii***

Raquel Gómez Sánchez^{a,*}, Elena del Valle Sánchez^b, Patricia Sanz Aznar^a, Ruth García Romero^{c,d}

^a Servicio de Pediatría, Hospital Universitario Miguel Servet, Zaragoza, Spain

^b Anatomía Patológica, Hospital Universitario Miguel Servet, Zaragoza, Spain

^c Gastroenterología Pediátrica, Hospital Universitario Miguel Servet, Zaragoza, Spain

^d Instituto de Investigación Sanitaria de Aragón (IIS Aragón), Zaragoza, Spain




Figure 1 Multiple nodules in the antrum and lesser curvature of the stomach corresponding to homogeneous granular-type lateral-spreading tumors (LSTs).

Chronic gastritis is the most common clinical manifestation caused by *Helicobacter pylori*. There are other, less prevalent *Helicobacter* species that cause gastrointestinal disease, such as *Helicobacter heilmannii*, which is associated with nodular antral gastritis.^{1,2}

DOI of original article:

<https://doi.org/10.1016/j.anpedi.2025.503928>

* Corresponding author.

E-mail address: raquelgs198@gmail.com (R. Gómez Sánchez).

We present the case of a child aged 3 years who had multiple food allergies and lived in a rural area in close contact with cats, with a reported history of difficulty swallowing solids, epigastric pain and frequent vomiting. Blood tests revealed vitamin B₁₂ deficiency and iron-deficiency anemia refractory to treatment. Endoscopy allowed visualization of antral mucosa nodularity with a polypoid appearance (Fig. 1), and the histological examination revealed the presence of spiral-shaped bacteria compatible with *H. heilmannii* (Figs. 2 and 3); culture of the biopsy specimen was not performed. After diagnosis, the patient received triple




Figure 2 Active chronic follicular gastritis in the antral mucosa (hematoxylin-eosin, original magnification $\times 10$).




Figure 3 Presence of *Helicobacter*-like spiral-shaped bacteria, compatible with *H. heilmannii* (Warthin-Starry, original magnification $\times 40$).

therapy (omeprazole, amoxicillin and metronidazole) for 14 days, which achieved resolution of symptoms.

H. heilmannii is a bacterium that spreads through zoonotic transmission that is rare in our area and is strongly associated with pets.³ Although it is morphologically different from *H. pylori*, endoscopic and histologic findings may be very similar in both species. We ought to underscore the need of additional microbiological tests in cases of non-*pylori* *Helicobacter* infection to improve the management of affected patients.^{2,3} Conventional *H. pylori* treatment can achieve eradication of *H. heilmannii*.³

References

1. Mention K, Michaud L, Guimber D, Martin De Lasalle E, Vincent P, Turck D, et al. Characteristics and prevalence of *Helicobacter heilmannii* infection in children undergoing upper gastrointestinal endoscopy. *J Pediatr Gastroenterol Nutr.* 1999;29:533–9, <http://dx.doi.org/10.1097/00005176-199911000-00012>.
2. Mohammadi M, Talebi Bezmin Abadi A, Rahimi F, Forootan M. *Helicobacter heilmannii* colonization is associated with high risk for gastritis. *Arch Med Res.* 2019;50:423–7, <http://dx.doi.org/10.1016/j.arcmed.2019.11.001>.
3. Chang D, Gluchowski NL, Abu Alfa AK, Goldsmith JD, Redston M, Bonilla S. Clinical presentation and outcomes of *Helicobacter heilmannii* gastritis in children in the New England region of the United States. *J Pediatr Gastroenterol Nutr.* 2024;78:204–10, <http://dx.doi.org/10.1002/jpn3.12058>.