

## IMAGES IN PAEDIATRICS

## Internal iliac artery pseudoaneurism after cardiac interventional procedure. Is conservative management feasible?

### Seudoaneurisma de arteria ilíaca interna tras cateterismo cardiaco intervencionista. ¿Es factible el manejo conservador?

Luis Fernandez Gonzalez<sup>a,\*</sup>, Roberto Blanco Mata<sup>a</sup>, Josune Arriola Meabe<sup>a</sup>, Javier Ayala Curiel<sup>b</sup>

<sup>a</sup> *Cardiología Intervencionista, Hospital Universitario de Cruces, Barakaldo, Vizcaya, Spain*

<sup>b</sup> *Cardiología Pediátrica, Hospital Universitario de Cruces, Barakaldo, Vizcaya, Spain*

We present the case of a boy aged 15 months with a prenatal diagnosis of pulmonary atresia with intact ventricular septum who underwent pulmonary valve perforation and balloon valvuloplasty with an 8mm balloon at 9 days post birth (Fig. 1A–D; Appendix B, videos 1–3). At 30 days, due to persistent desaturation, the decision was made to perform percutaneous ductal stenting, since this was less invasive compared to other techniques used to increase pulmonary flow. The procedure, guided by ultrasound, consisted in the delivery, via the left femoral artery and using a 5F introducer, of a drug-eluting stent (Onyx Frontier, Medtronic, USA) measuring 4.5 × 12 mm to the ductus arteriosus (Fig. 1E and F; Appendix B, videos 4 and 5). Five days after the procedure, the patient experienced vomiting, and the abdominal ultrasound scan used for initial assessment revealed a mass in the left groin, with subsequent Doppler and angio-CT


scans allowing visualization of a pseudoaneurysm at the level of the left common iliac artery (Fig. 2figureA–F).

Due to the risks involved in surgical intervention and the scarcity of the evidence on endovascular treatment in newborns, currently limited to isolated case reports,<sup>1,2</sup> the patient was managed conservatively with close clinical and radiological monitoring. Fifteen months later, the patient remains asymptomatic and has started to stand and walk without any problems associated with the pseudoaneurysm, which has exhibited progressive reduction in follow-up imaging (Fig. 3A and B). We present this case to demonstrate the feasibility of the conservative management of vascular complications in pediatric patients, an approach that requires very close clinical and radiological monitoring in experienced centers.


DOI of original article: <https://doi.org/10.1016/j.anpedi.2025.503838>

\* Corresponding author.

E-mail address: [luisfg82@hotmail.com](mailto:luisfg82@hotmail.com) (L. Fernandez Gonzalez).



**Figure 1** (A) and (B) Right ventriculogram, anteroposterior and lateral views showing pulmonary valve atresia (7.5 mm ring). (C) Pulmonary valvuloplasty performed with a TYSHAK MINI® (NuMED) 8 × 20 mm balloon catheter advanced on a 0.0014 coronary guidewire, which traversed the ductus arteriosus to the aorta supported by a gooseneck snare. (D) Angiographic confirmation of preserved pulmonary flow after valvuloplasty. (E) Lateral aortogram showing patent ductus arteriosus and filling of right chambers. (F) Aortogram following stent placement. PV, pulmonary valve; RV, right ventricle.



**Figure 2** (A)–(C) Doppler ultrasound: longitudinal (20 mm) and transverse (14 × 10 mm) views of the pseudoaneurysm with turbulent flow within. (D)–(F) Angio-CT images confirming the presence of a pseudoaneurysm arising from the left internal iliac artery and patency of distal vessels.




Figure 3 Follow-up ultrasound scan at 15 months showing a significant reduction in size (4 mm).

### Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.anpedi.2025.503838>.

### References

1. Pezzullo JA, Wallach MT. Successful percutaneous thrombin injection of a brachial artery pseudoaneurysm in a neonate. *AJR Am J Roentgenol.* 2002;178:244–5.
2. Cho MJ, Jeon UB, Choo KS, Lee HD. Percutaneous ultrasound-guided thrombin injection is effective even in infants with external iliac artery pseudoaneurysms. *Korean J Pediatr.* 2014;57:199–201.