



ORIGINAL ARTICLE

Compassion satisfaction, secondary traumatic stress and burnout: quality of working life in the field of pediatrics in Spain



María José Peláez Cantero^a, Iñigo Noriega Echevarría^b,
Juan Pablo García-Iñiguez^{c,*}, Felipe Verjano Sánchez^d, Ester Barrios Miras^e,
Inés del Río Pastoriza^f, María Jesús Alijas Merillas^g, Francisco Moreno Madrid^h,
Isolina Riaño Galánⁱ, José Antonio Salinas Sanz^j, en representación del Comité de
Bioética de la Asociación Española de Pediatría

^a Unidad de crónicos complejos y cuidados paliativos pediátricos en el Hospital Regional Universitario de Málaga, Spain

^b Hospital Universitario del Niño Jesús, Madrid, Universidad Internacional de La Rioja, La Rioja, Spain

^c Unidad Cuidados Intensivos Pediátricos, Hospital Universitario Miguel Servet, Zaragoza, Spain

^d Área Integrada de Pediatría y Neonatología, Hospital Universitario Costa del Sol, Marbella, Spain

^e Centro de Salud Mejorada del Campo, Madrid, Spain

^f Centro de Salud Arcade, Pontevedra, Spain

^g Clínica Universitaria de Navarra, Navarra, Spain

^h Servicio Pediatría, Hospital Clínico San Cecilio, Granada, Spain

ⁱ Área Gestión Clínica Pediatría, Hospital Universitario Central, Asturias, Spain

^j Servicio Onco-Hematología Infantil, Hospital Universitario Son Espases, Palma, Spain

Received 4 December 2024; accepted 4 August 2025

Available online 11 November 2025

KEYWORDS

Burnout;
Compassion
satisfaction;
Compassion fatigue;
Quality of working
life;
Secondary traumatic
stress

Abstract

Objective: To determine the prevalence of compassion satisfaction (CS), secondary traumatic stress (STS) and job exhaustion or burnout (BO) in medical professionals specialized in pediatrics at the national level in Spain and determine which demographic and work-related factors affect their development.

Methods: We conducted a cross-sectional study in pediatricians by means of questionnaires sent by the Spanish Association of Pediatrics (AEP) to its members, which were completed online and anonymously. We collected data on demographic variables, professional category (medical intern/resident [MIR] in pediatrics or pediatrician), main care setting and type of employment, specific field within pediatrics, main field of work, duration of work experience in general and time in current position. Care settings were further categorized into three groups: out-of-hospital, low-volume hospital and high-volume hospital (neonatology, intensive care, palliative

DOI of original article: <https://doi.org/10.1016/j.anpedi.2025.504012>

* Corresponding author.

E-mail address: dr_jpablo@hotmail.com (J.P. García-Iñiguez).

care, oncology and emergency care). Participants completed the Spanish adaptation of the Professional Quality of Life Scale (ProQOL) version 5 (*Escala de Calidad de Vida Profesional*) to assess three domains—CS, STS and BO—in relation to the past 30 days.

Results: We obtained a total of 1112 responses from pediatricians. Female respondents amounted to 78.9% of the sample. The distribution by care setting was 35.6% primary care, 34.9% low-volume hospital settings and 29.5% high-volume hospital settings. Most participants scored in the midrange of the three subscales of the ProQOL questionnaire: compassion satisfaction 60.7% (95% CI, 57.8–63.5), burnout 88.8% (95% CI, 86.8–90.5) and secondary traumatic stress 77.2% (95% CI, 74.7–79.6). Women scored significantly higher in the compassion fatigue subscales (BO and STS), while older age was associated with a linear increase in CS and an exponential decrease in STS. Permanent staff scored higher in CS and lower in BO and STS. We found a higher CS score in association with high-load hospital specialties and a higher BO score in association with low-load hospital specialties.

Conclusions: The surveyed sample of Spanish pediatricians showed significant levels of compassion fatigue and secondary traumatic stress, with greater impact among younger and less experienced professionals, temporary workers and female doctors, highlighting the need for further study and targeted educational interventions.

© 2025 Asociación Española de Pediatría. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Satisfacción por
compassion;
Fatiga por
compassion;
Estrés traumático
secundario;
Burnout;
Agotamiento laboral;
Calidad de vida
profesional

Satisfacción por compasión, estrés traumático secundario y agotamiento laboral: calidad de vida profesional en la Pediatría española

Resumen

Objetivos: Determinar la prevalencia de satisfacción por compasión (SC), estrés traumático secundario (SET) y burnout o agotamiento laboral (BO) en los profesionales médicos con especialidad de Pediatría a nivel nacional en España y determinar qué componentes demográficos y relacionados con el trabajo afectan a su desarrollo.

Método: Estudio transversal entre pediatras mediante encuestas enviadas desde la Asociación Española de Pediatría (AEP) a sus miembros, rellenadas virtualmente y en formato anónimo. Se recogieron variables demográficas, así como datos relativos a su situación profesional: pediatra en formación sanitaria especializada (MIR) o especialista en pediatría, ámbito principal de trabajo, tipo de relación laboral, área específica dentro de la pediatría, ámbito fundamental de trabajo, tiempo trabajado y tiempo en el puesto actual de trabajo. Las áreas específicas fueron categorizadas a su vez en tres grupos: extrahospitalarias, hospitalarias de baja carga y hospitalarias de alta carga (neonatología, cuidados intensivos, cuidados paliativos, oncología y urgencias). Los pediatras autocumplimentaron el cuestionario *Professional Quality of Life Scale* (ProQOL) en su versión 5 en castellano (*Escala de Calidad de Vida Profesional*) aplicado a los 30 días previos a su realización para evaluar al profesional en los tres dominios: SC, SET y BO.

Resultados: Se obtuvieron 1112 respuestas de pediatras (el 78,9% mujeres). Por áreas específicas los participantes se distribuyeron en Atención primaria 35,6%, Hospitalaria de baja carga 34,9% y Hospitalaria de alta carga 29,5%. La mayoría de los participantes se situaron en el rango medio de las tres subescalas del cuestionario ProQOL: satisfacción por compasión (SC) 60,7% (IC95%: 57,8–63,5), burnout (BO) 88,8% (IC95%: 86,8–90,5) y estrés traumático secundario (SET) 77,2% (IC95%: 74,7–79,6). Se observó una puntuación significativamente mayor en las subescalas de FC (BO y SET) para mujeres, mientras que una mayor edad se relacionó con un aumento lineal de SC y un descenso exponencial de SET. El personal fijo presentó mayor puntuación en SC y menor en BO y SET. Se observó una mayor puntuación en SC para especialidades hospitalarias de alta carga y mayor puntuación de BO en las de baja carga.

Conclusiones: La población de pediatras españoles encuestados presentó niveles de afectación relevantes de FC y TS, más importantes en profesionales de menor edad, mujeres y en profesionales temporales, siendo necesario el estudio y abordaje formativo de estos aspectos en estos grupos en particular.

© 2025 Asociación Española de Pediatría. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la CC BY-NC-ND licencia (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

At present, health care professionals, especially doctors, are facing challenging times. Health care organizations demand that professionals be committed, quick, adaptable, resilient and engaged in improving the quality of institutions to develop more efficient care delivery models and increase productivity¹: that they do more with less. In addition, the ongoing evaluation of patient satisfaction and assessment of health care quality with a greater focus on efficiency than on effectiveness can add to the pressure experienced by providers.

This ever-growing pressure has led many physicians to feel exhausted and unmotivated, weakening the bond between the medical profession and society. As a result, today we face a nationwide shortage of health care professionals, particularly in certain specialties.^{2–4} Multiple causes contribute to this phenomenon, involving various professional and social factors (social expectations, new models of care, progressive bureaucratization).⁵ Recent studies show that provider burnout has a negative impact on quality of care and patient safety and satisfaction.^{6,7} Furthermore, there is also evidence that this source of distress in physicians is associated with inadequate prescribing practices, ordering of unnecessary tests, an increased risk of malpractice lawsuits and lack of adherence of patients to the provider recommendations.^{7–11} A recent study estimated the prevalence of burnout in pediatricians at up to 36%, warranting a more in-depth study of this issue.⁵

In consequence, monitoring these issues and knowing the current situation has become an imperative.¹² To this end, Stamm conceived of a scale to measure the perceived quality of one's work, developing the Professional Quality of Life (ProQOL) tool. Its purpose is screening for issues in professional quality of life (pQoL), which has two key dimensions: compassion satisfaction (CS) and compassion fatigue (CF). In turn, compassion fatigue is composed of two aspects: secondary traumatic stress (STS) and burnout (BO).

The aim of our study was to determine the prevalence of CS, STS and BO among physicians specialized in pediatrics in Spain and delve into the demographic and work-related factors associated with their development. The nature of the study was chiefly descriptive, through an exploratory analysis of the data, with no predefined working hypothesis, given the absence of previous similar studies in the field of pediatrics in Spain.

Material and methods

We conducted a cross-sectional study by means of self-report questionnaires distributed through the Asociación Española de Pediatría (AEP, Spanish Association of Pediatrics) to its members between February and March 2024, which were completed online on an anonymous basis. We collected data on demographic variables (sex/gender, age) and work-related variables, including whether the respondent was training as a medical resident/intern (MIR) in pediatrics or was an accredited pediatrician/pediatric specialist, the care setting and the type of institution (primary

care center vs hospital) where the respondent worked most of the time, the type of employment contract (temporary/permanent), specific field within pediatrics, total years of experience and years worked in current position. Specific care settings were also categorized into three groups: out-of-hospital/primary care, low-acuity hospital-based care and high-acuity hospital-based care. The latter category included neonatal care, intensive care, palliative care, oncology and emergency care, based on the criteria of the researchers leading the project.

We also collected data corresponding to the Spanish adaptation of the ProQOL version 5, using the official manual to interpret the results, except in cases in which we opted to not standardize the results to facilitate their understanding.¹³ The scale comprises 30 items rated on a 5-point Likert scale (from 0 [*never*] to 5 [*very often*]) that apply to the 30 days before completion to assess the following domains in the provider: CS, STS and BO. The compassion satisfaction subscale assesses the pleasure and satisfaction that providers derive from contributing to the wellbeing of patients and their families.¹⁴ On the other hand, the two subscales for compassion fatigue, STS and BO, differ both in nature and in their underlying causes. Secondary traumatic stress refers to the stress derived from the indirect exposure to the trauma experienced by others, usually through listening to the traumatic experiences reported by the patients, and therefore stems from an excess of empathy toward the traumatized patients,^{15,16} while BO is a state of emotional exhaustion, depersonalization and low personal fulfilment due to chronic work-related stressors, such as excessive workloads, interpersonal conflict and the lack of resources or support in the workplace.^{17,18}

The instrument yields a score for each of these three subscales, which allow categorization of respondents into 3 groups: *low*, *medium* or *high*. The boundary between low/medium and medium/high scores correspond approximately to the 25th and 75th percentile in the population (23 and 43 points, respectively). While *high* scores in CS correspond to a higher pQoL, high scores in the BO and STS are indicative of a poorer pQoL. Different studies have found a good content and construct validity as well as adequate reliability for the ProQOL^{13,19} (Table 1).

We summarize quantitative data as mean and standard deviation and categorical data as percentages. We also conducted an exploratory analysis with bivariate tests based on the epidemiological variables of sex/gender and age, comparing MIR trainees versus the accredited pediatrician categories noted above, type of employment contract and years spent in the current position (excluding MIR trainees from the two latter variables). Comparisons were made with the Student *t* test or analysis of variance (ANOVA) depending on the number of categories. In the case of ordinal categories, we also analyzed whether they followed a linear or exponential trend. In addition, we used backward stepwise logistic regression to analyze the association of each of the subscales with the study variables. The statistical analysis was performed with the Stata software, version 16.1.

Last of all, the study was exempt from explicit request of informed consent given that questionnaires were self-administered on a voluntary basis.

Table 1 Components of the Professional Quality of Life (ProQOL) scale.

	ProQOL score	Professional quality of life
Compassion satisfaction	High	Good
	Low	Poor
Compassion fatigue		
Secondary traumatic stress	High	Poor
	Low	Good
Burnout	High	Poor
	Low	Good

Results

We received a total of 1112 responses, with a response rate of 8.2%. Table 2 presents the main epidemiological results. There was a clear predominance of female respondents and a relatively uniform age distribution, and a majority of respondents held permanent positions (62.5%).

Table 3 presents the overall results of the ProQOL. Most participants scored in the medium range for the three subscales, with very small percentages in the groups with low CS or high BO or STS.

Table 4 summarizes the results of the exploratory comparative analysis. In the case of sex/gender, for mathematical reasons, we excluded the response corresponding to the "other" gender category, which scored in the medium range for every variable. We found significantly higher scores in the CF subscales (BO and STS) among female respondents. Increasing age was associated with a linear increase in CS and an exponential decrease in STS.

As regards work-related variables, MIR trainees scored lower in CS and higher in the two CF scales, while an increasing number of years in the current position was associated with a linear increase in CS and a linear decrease in BO, with significant differences in STS but without a discernible ordinal pattern. In the analysis by type of employment, we eventually chose to exclude respondents who were retired or self-employed due to the small size of the subset compared to those who held permanent and temporary positions. Permanent staff scored higher in CS and lower in BO and STS. We also found differences in the scores of the three subscales based on the pediatric subspecialty categories. Although the methods did not allow for differentiation among each of the different subspecialty categories, we found higher CS scores among respondents working in high-acuity hospital care settings and higher BO scores among those working in low-acuity hospital care settings, while the difference could not be evaluated for STS.

When we fitted linear regression models for each of the primary study variables, in the case of CS, the factors that were included in the final model were working in a high-acuity subspecialty (coefficient, 1.4), having held the current position for 20 or more years (coefficient, 1.2) and greater age, with the correlation coefficient increasing progressively with increasing age (coefficient, 1.0 for age 36–45 years, 2.0 for age 46–55 years and 3.0 for age >56 years). In the BO model, the associated variables were female sex (coefficient, 1.0), age, although there was only a significant difference between the oldest age group compared to all others (coefficient, −2.0), temporary employment

Table 2 Demographic and work-related characteristics of the pediatricians who participated in the study.

	Percentage
Sex/gender	
Female	78.9
Male	21.0
Other	0.1
Age	
25–35	27.5
36–45	28.3
46–55	21.5
>56	22.7
Employment (n = 1098)	
MIR trainee	11.4
Temporary	24.0
Permanent	62.5
Self-employed	1.7
Retired	0.5
Care setting (excluding MIR)	
Primary care	43.5
Hospital	56.5
Subspecialty category (excluding MIR; n = 891)	
Primary care	35.6
Low-acuity hospital setting	34.9
High-acuity hospital setting	29.5
Years of experience (excluding MIR; n = 967)	
<5 years	12.7
5–10 years	16.1
10–15 years	17.9
16–20 years	13.0
>20 years	40.2
Years in current position (excluding MIR; n = 959)	
<5 years	34.1
5–10 years	22.1
10–15 years	14.2
16–20 years	11.6
>20 years	18.0

For categories in which there are missing data, we present the number of participants included in the analysis (n). For the variables for which MIR trainees were excluded, we only show the n value if there were missing data.

Abbreviation: MIR, medical intern-resident trainee.

(coefficient, 1.5) and working in a low-acuity hospital-based pediatric care setting (coefficient, 0.9). For the TS subscale, the factors identified as determinants in the model were female sex (coefficient, 2.1), working in a low-acuity hos-

Table 3 Results obtained in the total sample of pediatricians for the compassion satisfaction, burnout and secondary traumatic stress subscales of the ProQOL scale for analysis of professional quality of life, expressed as a mean or percentage and 95% confidence interval.

	Mean or percentage (95% CI)
Compassion satisfaction (CS)	38.1 (37.7–38.4)
Low	0.6% (0.3–0.13)
Medium	60.7% (57.8–63.5)
High	38.7% (35.8–41.6)
Burnout (BO)	29.6 (29.2–29.9)
Low	8.1% (6.6–9.9)
Medium	88.8% (86.8–90.5)
High	3.1% (22.7–43.5)
Secondary traumatic stress (STS)	26.2 (25.9–26.6)
Low	21.5% (19.2–24.0)
Medium	77.2% (74.7–79.6)
High	1.3% (0.7–2.1)

pital setting (coefficient, 1.0) and temporary employment (coefficient, 0.9).

Discussion

Our study, conducted through the self-administration by pediatric providers of the ProQOL questionnaire for assessment of professional quality of life, found significant levels of CF, measured in terms of both BO and STS, among the pediatricians working in Spain that submitted responses. Compassion fatigue was greater in providers who were younger, female or held temporary positions.

The findings of our study in Spanish pediatricians were similar to those reported in other populations, evincing the need to address the issue of CF in our country. We identified several factors potentially associated with a poorer pQoL, chief among them age, temporary employment and female sex. The scores in our sample for all three subscales were similar to those reported in a meta-analysis that included data from 11 countries²⁴ and other studies published in countries such as Nepal,²⁵ United States²⁶ or Vietnam.²⁷

The potential protective factors for pQoL identified in our study were greater age and longer work experience, which

Table 4 Comparison of the results obtained in the subscales of the ProQOL based on sex/gender, age, professional category (MIR vs specialist), subspecialty category/care setting, type of employment and time in current position (MIR trainees excluded from the last three).

	CS		BO		STS
		P value		P value	P value
Sex/gender		.1		< .01*	< .01*
Female	37.9 (0.2)		29.9 (0.2)		26.6 (0.2)
Male	38.6 (0.4)		28.3 (0.4)		24.7 (0.4)
Age		.047*		.06	< .01*
25–35	36.9 (0.4)		31.1 (0.3)		26.5 (0.3)
36–45	37.7 (0.3)		30.0 (0.3)		26.4 (0.3)
46–55	38.8 (0.4)		29.4 (0.4)		26.4 (0.4)
> 56	40.3 (0.4)		27.2 (0.4)		25.5 (0.4)
Physician category		< .01*		< .01*	< .03*
MIR	35.0 (0.5)		32.3 (0.5)		27.3 (0.5)
Specialist in pediatrics	38.5 (0.2)		29.2 (0.2)		26.1 (0.2)
Employment status		< .01*		< .01*	.03*
Temporary	37.2 (0.3)		30.7 (0.3)		26.8 (0.4)
Permanent	38.9 (0.2)		28.7 (0.2)		25.9 (0.2)
Care setting		< .01*		< .01*	< .01*
Primary Care	38.0 (0.3)		28.8 (0.3)		25.5 (0.3)
Low-acuity hospital setting	38.0 (0.3)		30.0 (0.3)		26.7 (0.3)
High-acuity hospital setting	39.3 (0.4)		28.9 (0.3)		26.0 (0.4)
Years in current position		.04*		.02*	< .01*
<5 years	37.5 (0.3)		29.9 (0.3)		26.1 (0.3)
5–10 years	37.7 (0.4)		29.6 (0.4)		25.9 (0.4)
10–15 years	38.9 (0.5)		28.9 (0.5)		26.2 (0.5)
16–20 years	39.4 (0.5)		29.2 (0.5)		26.4 (0.6)
>20 years	40.6 (0.4)		27.4 (0.4)		25.8 (0.4)

Scores expressed as mean (SD) for each group. *Statistically significant.

Abbreviations: BO, burnout; CS, compassion satisfaction; MIR, medical intern-resident trainee; ProQOL, Professional Quality of Life scale; STS, secondary traumatic stress.

was consistent with a large volume of evidence showing that older and more experienced providers exhibit significantly greater CS and lesser BO,^{25,27–30} as MIR trainees scored lower in CS and higher in the two CF subscales and, as the number of years in the current position increased, there was a linear increase in CS and a linear decrease in BO, in addition to significant differences in STS. A possible interpretation is that older and more experienced providers, due to their greater exposure to the difficulties and challenges faced by patients, may become more adaptable and be better equipped to manage their own stress and exhaustion.^{27,31} In contrast, younger and more inexperienced workers may not be mentally prepared to face the extreme situations that patients experience on a daily basis.

One of the salient findings of our study was the association of provider sex with CF. A recent meta-analysis of studies conducted among physicians in Spain did not identify differences in relation to sex/gender, with the authors suggesting, nonetheless, that it is important to consider the gender perspective.^{32,33} Our study, conducted in a group with a large proportion of female providers (pediatric care), supports this view.

In respect of the pediatric subspecialties, we found higher CS scores in providers working in high-acuity hospital-based settings (neonatology, intensive care, palliative care, oncology and emergency care) and higher BO scores in low-acuity hospital-based settings (all other). The increased exposure of providers to complex situations in high-acuity specialties, involving considerable suffering in children and their families, may be associated with increased development of adequate protective strategies to prevent BO and STS in providers while promoting CS. Previous studies have found differences in relation to acuity, showing a lower prevalence of BO in surgery departments compared to other specialties,^{34,35} a higher prevalence of BO and STS in intensive care units,^{24,34} and a higher prevalence of CS and lower prevalence of BO in palliative care.³⁶

However, we did not find scientific evidence on the association between the type of employment contract and pQoL, while one of the findings in our study was that providers with permanent positions scored higher on CS and lower on BO and STS. Job insecurity may be an organizational barrier hindering compassionate care.

There are several limitations to this study. First of all, we need to highlight the low response rate relative to the total number of AEP members (1112 responses out of a total of 13 584 members), although the number of submitted responses can be considered high for this type of study. There is also a risk of selection bias, as pediatricians who are more affected could be more likely to respond. Other factors, such as time constraints or the ease of completing the questionnaire online, could also lead to differential participation among the population of pediatricians. In addition, we did not study other variables that have been found to be associated with pQoL, such as income^{27,37} or marital status.^{30,34,38} In any case, we did not establish a working hypothesis and the performance of multiple comparisons could have led us to find spurious associations. Even the results of the logistic regression analysis should be interpreted with caution, given the sampling method. Future studies should randomly recruit pediatricians or even quota sampling to allow a more reliable analysis of the impact of various factors on the out-

comes of interest. Among the strengths of the study, we ought to highlight its focus on pediatric care and the assessment of CF in pediatricians who practice in Spain by means of a questionnaire validated for use in the Spanish population. The pQoL of pediatricians is an aspect that needs to be studied in detail to protect and promote current and future health care quality.

In conclusion, our study found CF, STS and CS levels in a cohort of pediatricians in Spain that were similar to those reported in studies conducted in other countries, evincing a poorer pQoL among providers of younger age, female sex or with temporary positions. Future studies and interventions should consider specific factors and opportunities for improvement, specifically including coping strategies appropriate for the needs of pediatricians in training and a gender perspective.

Ethical considerations

The study adhered to good clinical practice and ethical principles for medical research involving human subjects established by the World Medical Association in the Declaration of Helsinki and its subsequent amendments through the last version from 2024, as well as the Convention on Human Rights and Biomedicine.^{20,21} On distributing the questionnaire, participants were informed that the information collected in the survey would be solely and exclusively used for the stated purposes, safeguarding the anonymity of respondents who completed it on a voluntary basis, in addition to the confidentiality of the collected data in adherence to Regulation (EU) 2016/679 and Organic Law 3/2018 of 5 December on the Protection of Personal Data and Guarantee of Digital Rights.^{22,23}

Declaration of competing interest

The authors have no conflicts of interest to declare.

Acknowledgments

We thank the Asociación Española de Pediatría for enabling the distribution of the questionnaire among its members, as well as the members themselves for the time they devoted to responding.

References

- Swensen S, Kabacell A, Shanafelt T. Physician-organization collaboration reduces physician burnout and promotes engagement: the mayo clinic experience. *J Healthc Manag Am Coll Healthc Exec.* 2016;61(2):105–27.
- SÁNCHEZ LG. RTVE.es. 2022 [citado 10 de junio de 2024]. Sobrecargados, infravalorados y desprestigiados: por qué muchos MIR ya no quieren ser médicos de familia. Disponible en: <https://www.rtve.es/noticias/20220525/medicos-familia-atencion-primaria-espana/2350895.shtml>.
- Shanafelt TD, Boone S, Tan L, Dyrbye LN, Sotile W, Satele D, et al. Burnout and satisfaction with work-life balance among US physicians relative to the general US population. *Arch Intern Med.* 2012;172(18):1377.

4. Shanafelt TD, Hasan O, Dyrbye LN, Sinsky C, Satele D, Sloan J, et al. Changes in burnout and satisfaction with work-life balance in physicians and the general US working population between 2011 and 2014. *Mayo Clin Proc.* 2015;90(12):1600–13.
5. Peirau M, Esquerda M, Carme Gabarrell, Pifarre J. Veinte años después: prevalencia y evolución del burnout en Pediatría, de 1998-1999 a 2018-2019. *Rev Pediatr Aten Primaria* [Internet]. 2021;23(91):253–60. Epub 06-Feb-2023. Available from: http://scielo.isciii.es/scielo.php?script=sci_arttext&pid=S1139-76322021000300004&lng=es
6. Association of Resident Fatigue and Distress With Perceived Medical Errors | Depressive Disorders | JAMA | JAMA Network [Internet]. [access 5 June 2024]. Available from: <https://jamanetwork.com/journals/jama/article-abstract/184625>.
7. Balch CM, Oreskovich MR, Dyrbye LN, Colaiano JM, Satele DV, Sloan JA, et al. Personal consequences of malpractice lawsuits on American surgeons. *J Am Coll Surg.* 2011;213(5):657–67.
8. Melville A. Job satisfaction in general practice: implications for prescribing. *Soc Sci Med [Med Psychol Med Sociol]*. 1980;14A(6):495–9.
9. DiMatteo MR, Sherbourne CD, Hays RD, Ordway L, Kravitz RL, McGlynn EA, et al. Physicians' characteristics influence patients' adherence to medical treatment: Results from the Medical Outcomes Study. *Health Psychol.* 1993;12(2):93–102.
10. Grol R, Mokkink H, Smits A, van Eijk J, Beek M, Mesker P, et al. Work satisfaction of general practitioners and the quality of patient care. *Fam Pract.* 1985;2(3):128–35.
11. Jones JW, Barge BN, Steffy BD, Fay LM, Kunz LK, Wuebker LJ. Stress and medical malpractice: organizational risk assessment and intervention. *J Appl Psychol.* 1988;73(4):727–35.
12. Wallace JE, Lemaire JB, Ghali WA. Physician wellness: a missing quality indicator. *The Lancet.* 2009;374(9702):1714–21.
13. B. Hudnall Stamm, 2009-2012. Professional Quality of Life: Compassion Satisfaction and Fatigue Version 5 (ProQOL).
14. Sacco TL, Copel LC. Compassion satisfaction: a concept analysis in nursing. *Nurs Forum (Auckl)*. 2018;53(1):76–83.
15. Figley CR. *Compassion Fatigue*. New York: Taylor & Francis Group, LLC; 1995.
16. Sabo BM. Compassion fatigue and nursing work: can we accurately capture the consequences of caring work? *Int J Nurs Pract.* 2006;12(3):136–42.
17. Edú-Valsania S, Laguía A, Moriano JA. Burnout: a review of theory and measurement. *Int J Environ Res Public Health.* 2022;19(3):1780.
18. Montero-Marín J [access 11 June 2024]. Available from: <https://www.semanticscholar.org/paper/The-burnout-syndrome-and-its-various-clinical-a-for-Montero-Mar%C3%ADn/a32f3fbd9fbc2dfadfb07990fc576880ebe8ba1>, 2016.
19. Singh J, Karanika-Murray M, Baguley T, Hudson J. A psychometric evaluation of Professional Quality of Life Scale Version 5 (ProQOL 5) in a UK-based sample of allied mental health professionals. *Curr Psychol.* 2024;43(25):21615–29.
20. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human participants. *JAMA.* 2025;333(1):71.
21. Council of Europe. In: Convention for the protection of Human Rights and Dignity of the Human Being with regard to the Application of Biology and Medicine: Convention on Human Rights and Biomedicine (ETS No. 164) [Internet]; 1997 [access 1 January 2021]. Available from: <https://www.coe.int/en/web/conventions/full-list?module=treaty-detail&treatynum=164>
22. Comité de Ministros, Consejo de Europa. Recomendación Rec (2003) 24 del Comité de Ministros de los estados miembros sobre organización de cuidados paliativos. Available from: https://www.mscbs.gob.es/organizacion/sns/planCalidadSNS/pdf/excelencia/cuidadospaliativos-diabetes/CUIDADOS_PALIATIVOS/opsc_est6.pdf.pdf. Accessed 13 May 2021.
23. España. Ley Orgánica 3/2018, de 5 de diciembre, de Protección de Datos Personales y garantía de los derechos digitales. Boletín Oficial del Estado. 2018 Dic 6;(294):119788–119820. Available from: <https://www.boe.es/eli/es/lo/2018/12/05/3/con>.
24. Xie W, Chen L, Feng F, Okoli CTC, Tang P, Zeng L, et al. The prevalence of compassion satisfaction and compassion fatigue among nurses: a systematic review and meta-analysis. *Int J Nurs Stud.* 2021;120:103973.
25. Vaidya A, Karki S, Dhimal M, Gyanwali P, Baral D, Pandey A, et al. Professional quality of life among medical doctors working in Kathmandu: a descriptive cross-sectional study. *JNMA J Nepal Med Assoc.* 2020;58(231):900–4.
26. Hunsaker S, Chen HC, Maughan D, Heaston S. Factors that influence the development of compassion fatigue, burnout, and compassion satisfaction in emergency department nurses. *J Nurs Scholarsh Off Publ Sigma Theta Tau Int Honor Soc Nurs.* 2015;47(2):186–94.
27. Tran ANP, To QG, Huynh VAN, Le KM, To KG. Professional quality of life and its associated factors among Vietnamese doctors and nurses. *BMC Health Serv Res.* 2023;23(1):924.
28. El-Shafei DA, Abdelsalam AE, Hammam RAM, Elgohary H. Professional quality of life, wellness education, and coping strategies among emergency physicians. *Environ Sci Pollut Res Int.* 2018;25(9):9040–50.
29. ResearchGate [Internet]. [access 13 June 2024]. Available from: https://www.researchgate.net/publication/330708599_The_professional_quality_of_life_among_health-care_providers_and_its_related_factors/link/5c504596458515a4c7480cb1/download?tp=eyJjb250ZXh0Ijp7ImZpcnN0UGFnZSI6InB1YmVpY2F0aW9uIiwicGFZSI6InB1YmVpY2F0aW9uIn19.
30. Su PA, Lo MC, Wang CL, Yang PC, Chang CI, Huang MC, et al. The correlation between professional quality of life and mental health outcomes among hospital personnel during the Covid-19 pandemic in Taiwan. *J Multidiscip Healthc.* 2021;14:3485–95.
31. Peisah C, Latif E, Wilhelm K, Williams B. Secrets to psychological success: why older doctors might have lower psychological distress and burnout than younger doctors. *Aging Ment Health.* 2009;13(2):300–7.
32. Pujol-de Castro A, Valerio-Rao G, Vaquero-Cepeda P, Catalá-López F. Sexo/género en estudios de prevalencia del síndrome de burnout en médicos: análisis de metarregresión. *Gac Sanit* [Internet]. 2024;38:102404, <http://dx.doi.org/10.1016/j.gaceta.2024.102404>. Epub 06-Dic-2024. Available from: http://scielo.isciii.es/scielo.php?script=sci_arttext&pid=S0213-91112024000100801&lng=es
33. Pujol-de Castro A, Valerio-Rao G, Vaquero-Cepeda P, Catalá-López F. Prevalencia del síndrome de burnout en médicos que trabajan en España: revisión sistemática y metaanálisis. *Gac Sanit.* 2024;38:102384.
34. Gümüş E, Alan H, Taşkıran Eskici G, Eşkin Bacaksız F. Relationship between professional quality of life and work alienation among healthcare professionals. *Florence Nightingale J Nurs.* 2021;29(3):342–52.
35. Rotenstein LS, Torre M, Ramos MA, Rosales RC, Guille C, Sen S, et al. Prevalence of burnout among physicians: a systematic review. *JAMA.* 2018;320(11):1131–50.
36. O'Mahony S, Ziadni M, Hoerger M, Levine S, Baron A, Gerhart J. Compassion fatigue among palliative care clinicians: findings on personality factors and years of service. *Am J Hosp Palliat Care.* 2018;35(2):343–7.

37. Mangoulia P, Koukia E, Alevizopoulos G, Fildissis G, Katostaras T. Prevalence of secondary traumatic stress among psychiatric nurses in Greece. *Arch Psychiatr Nurs*. 2015;29(5):333–8.
38. Aslan H, Erci B, Pekince H. Relationship between compassion fatigue in nurses, and work-related stress and the meaning of life. *J Relig Health*. 2022;61(3):1848–60.