

Koolen de Vries syndrome: A challenge in clinical practice[☆]



Síndrome Koolen de Vries: un reto en la práctica clínica

Dear Editor:

Koolen-de Vries syndrome (KDVS, OMIM 610443) is a rare genetic disorder with an approximate prevalence of 1 per 16,000 live births, with no predominance of either sex. It is characterised by neonatal hypotonia, intellectual disability, dysmorphic features (high and broad forehead, long face, palpebral fissures with epicanthic folds, pear-shaped nose) that become attenuated over the years, and friendly behaviour. Central nervous system abnormalities are also found in 80% of cases, such as epilepsy (50%) and brain malformations (hydrocephalus and agenesis or dysgenesis

of the corpus callosum), congenital heart defects (valvulopathies and septal defects) in 40–63% depending on the series, and urogenital anomalies (cryptorchidism, hypospadias, vesicoureteral reflux, hydronephrosis, etc.) in up to 70%.¹

It was first described in 2006 as a recurrent microdeletion localised in chromosome 17q21.31, between 500 and 650 kb in length, that comprises up to five genes: *CRHR1*, *SPPL2C*, *MAPT*, *STH* and *KANSL1* (or *KIAA1267*), in addition to two putative genes, *MGC57346* and *C17orf69*. In most cases, these are *de novo* mutations. Although the role of these genes in the pathogenesis of this disease remains unclear, recent studies have demonstrated that both *KANSL1* haploinsufficiency or point mutations are sufficient to produce the disease.^{2–4} The most recently published case series (with 45 patients) described that while the phenotype is not significantly different based on whether there is a 17q21.31 microdeletion or a nonsense mutation of the *KANSL1* gene, there is wide phenotypic variability between individuals.⁴

Table 1 Case characteristics.

| | Case 1 | Case 2 | Case 3 | Case 4 |
|-------------------------|--|--|--|---|
| Presenting complaint | Language delay | Psychomotor delay | Epilepsy | Delayed postural control |
| Phenotype | Dolichocephaly, long face, low-set prominent ears and bulbous nose | Hypertelorism, broad nasal root, long philtrum, low-set antverted ears, broadly spaced nipples and mild pectus excavatum | Skull disproportionate to face, broad forehead and low-set ears | Long face, hypertelorism, epicanthic folds and low-set ears |
| Language | Monosyllabic | Absent No verbal language | Language delay | Poor, with difficulty in verbal expression |
| Intellectual disability | Yes | Yes | Yes | Yes |
| Behaviour | Friendly, but with difficulty in communicating with peers | Sociable | Friendly Hyperactivity | Affable Attention deficit |
| Epilepsy | No | Yes | Yes | No |
| Brain MRI | Hypogenesis of corpus callosum | Hypoplasia of posterior corpus callosum | Ventriculomegaly in the absence of hydrocephalus. Cortical and subcortical atrophy | Very mild cortical and subcortical atrophy |
| Karyotype | Normal | Normal | Normal | Normal |
| Fragile X Array CGH | Normal 1600 kb microdeletion in the 17q21.31 band affecting up to 13 genes, including <i>MAPT</i> and <i>KANSL1</i> | Normal Deletion of chromosome 17q21.31 (tested in a different facility) | Normal 621 kb microdeletion in 17q21.31 affecting the <i>CRHR1</i> , <i>IMPS</i> , <i>MAPT</i> , <i>STH</i> and <i>KANSL1</i> genes | Normal 695 kb microdeletion in 17q21.31 affecting the <i>C17orf69</i> , <i>IMP5</i> , <i>MAPT</i> , <i>STH</i> and <i>KANSL1</i> genes |
| Echocardiogram | Normal | Bicuspid aortic valve | Normal | Normal |
| Renal ultrasound | Normal | Normal | Normal | Normal |
| Other | | Short stature | | Father with inversion polymorphism |

☆ Please cite this article as: Moreno Samos M, Moreno Medinilla EE, Martínez Antón JL, Urda Cardona A. Síndrome Koolen de Vries: un reto en la práctica clínica. An Pediatr (Barc). 2017;86:162–164.

Table 2 Comparison of the most relevant characteristics found in the most recently published series and in our patients.

| | Koolen, 2008 (%) N=22 | Zollino, 2015 (%) N=32 | Koolen, 2016 (%) N=45 | Our series (%) N=4 |
|----------------------------------|--------------------------|---------------------------|--------------------------|-----------------------|
| Dysmorphic features | | | | |
| Long face | 74 | 75 | 75 | 50 |
| Bulbous nose | 82 | 93.7 | 88.3 | 50 |
| Prominent ears | 59 | — | 32.6 | 75 |
| Everted lower lip | — | 93.7 | 71.4 | 0 |
| Epicantiche fold | 68 | — | 52.3 | 50 |
| Macrocephaly | — | 40 | 14.3 | 25 |
| Neurologic features | | | | |
| Hypotonia | 96 | 100 | 86.4 | 50 |
| Intellectual disability | — | 90 | 100 | 100 |
| Motor delay | 100 | — | 97.3 | 75 |
| Language delay | | 87 | 100 | 100 |
| Epilepsy | 50 | 50 | 48.9 | 50 |
| CNS structural malformations | — | 50 | 52.9 | 75 |
| Friendly/amiable behaviour | 89 | 95 | 88.7 | 100 |
| Musculoskeletal anomalies | | | | |
| Heart defects | 25 | 40 | 76.7 | 25 |
| Renal/urogenital anomalies | 27 | 35 | 38.6 | 25 |
| Short stature | 32 | 22 | 45.2 | 0 |
| | 18 | 37 | 35.3 | 25 |

A previously described predisposing factor is the inversion of chromosome 17 in one of the parents, which, while necessary, is not sufficient to produce the microdeletion, as it is a very common polymorphism that is found in up to 20% of the European population.¹ Only two cases of sibling recurrence have been reported in two independent families in which the mothers had a mosaicism for the chromosome 17 deletion, which could be a risk factor for recurrence, underscoring the importance of offering genetic counselling to parents of affected children.⁵

We present a series of four patients with KDVS (Table 1). All patients had some of the clinical features included in the 37 symptoms described by Koolen in the first published series; however, the diagnosis was made when expanded molecular testing was requested following an initial battery of diagnostic tests that were inconclusive. We present a table that summarises what we consider to be the most relevant clinical features described in the most recently published and broader series and those found in our patients (Table 2). Intellectual disability and facial dysmorphisms are the most frequently observed features, as was the case in our patients, although potential comorbidities need to be ruled out. Contrary to what has been reported in the literature, none of our patients presented with nephrourologic abnormalities.

Genetic testing in all cases identified mutations in one of the five genes that cause the disease. We ought to highlight the size of the mutation in case 1, which exceeded sizes reported to date.

Recently, the first case has been published of KDVS diagnosed prenatally through the detection of bilateral ventriculomegaly in the pregnancy ultrasound examination performed at 33 weeks' gestation, with confirmation of the microdeletion by means of array-based comparative

genomic hybridisation in a sample obtained by amniocentesis.⁶ This reinforces the importance of detecting the syndrome early, as it not only allows its aetiological diagnosis, but also genetic counselling for the family. Furthermore, considering that this is a fundamentally monogenic disorder, it is likely that it can benefit from pharmacologic treatment targeting the genome or proteome in the future if research is conducted on this field.

In conclusion, KDVS is a rare disease that must be included in the differential diagnosis of patients with unexplained intellectual disability with or without associated dysmorphic features or malformations. Considering the wide clinical variability that has been observed, we believe that genetic hybridisation techniques must be included at the level of brain MRI in the diagnostic evaluation of these patients.

References

1. Koolen DA, Sharp AJ, Hurst JA, Firth HV, Knight SJL, Goldenberg A, et al. Clinical and molecular delineation of the 17q21.31 microdeletion syndrome. *J Med Genet.* 2008;45:710–20.
2. Koolen DA, Kramer JM, Neveling K, Nillesen WM, Moore-Barton HL, Elmslie FV, et al. Mutations in the chromatin modifier gene *KANSL1* cause the 17q21.31 microdeletion syndrome. *Nat Genet.* 2012;44:639–41.
3. Zollino M, Marangi G, Ponzi E, Orteschi D, Ricciardi S, Lattante S, et al. Intragenic *KANSL1* and chromosome 17q21.31 deletions: broadening the clinical spectrum and genotype–phenotype correlations in a large cohort of patients. *J Med Genet.* 2015;52:804–14.
4. Koolen DA, Pfundt R, Linda K, Beunders G, Veenstra-Knol HE, Conta JH, et al. The Koolen-de Vries syndrome: a phenotypic comparison of patients with a 17q21.31 microdeletion versus a *KANSL1* sequence variant. *Eur J Hum Genet.* 2016;24:652–9.

5. Koolen DA, Dupont J, de Leeuw N, Vissers LE, van den Heuvel SP, Bradbury A, et al. Two families with sibling recurrence of the 17q21.31 microdeletion syndrome due to low-grade mosaicism. *Eur J Hum Genet.* 2012;20:729–33.
6. Egloff M, Encha-Razavi F, Garel C, Bonnière-Darcy M, Millischer AE, Lapierre JM, et al. 17q21.31 microdeletion: brain anomalies leading to prenatal diagnosis. *Cytogenet Genome Res.* 2014;144:178–82.

María Moreno Samos, Esther Eugenia Moreno Medinilla*,
Jacinto Luis Martínez Antón, Antonio Urda Cardona

Sección de Neuropediatría, Servicio de Pediatría, Hospital Materno Infantil, Hospital Regional Universitario de Málaga, Málaga, Spain

* Corresponding author.

E-mail address: esthermoreno84@hotmail.com
(E.E. Moreno Medinilla).

2341-2879/

© 2016 Asociación Española de Pediatría. Published by Elsevier España, S.L.U. All rights reserved.