



IMAGES IN PAEDIATRICS

Bead-like lesion at perineal region**Lesión de aspecto arrosariado a nivel perineal**

Juan Pablo Velasco-Amador*, Israel Pérez-López, Ricardo Ruiz-Villaverde

*Unidad de Gestión Clínica de Dermatología Médico Quirúrgica y Venereología, Hospital Universitario San Cecilio, Granada, Spain*

Available online 1 March 2024

A boy aged 3 years with an unremarkable history was referred to the paediatric dermatology clinic for assessment of perineal lesions present from birth. The physical examination revealed several small confluent cysts in a string of pearls arrangement over the midline of the penis extending from the distal end through the ventral raphe to the perianal region (Fig. 1). The lesions were asymptomatic and showed no signs of infection. An ultrasound scan of the kidney, urinary tract and the largest lesion was performed in which the lesion appeared hypoechoic with posterior enhancement.

Given the characteristics of the lesions and the absence of additional abnormalities on imaging, the patient received a final diagnosis of perineal median raphe cysts. Median raphe cysts are a rare congenital defect of the male genitals.¹ They are usually found between the urethral meatus and the perianal region, and the diagnosis is clinical.¹ They are considered a developmental defect resulting from the abnormal fusion of urethral folds into the genital tubercle in the foetus.² Most cases are asymptomatic, and the most frequent complication is overinfection.² Although this defect is benign, the first-line treatment is surgical correction with the aim of preventing complications.³



Figure 1 Cystic lesions in a pearl string arrangement along the midline of the raphe.

DOI of original article:

<https://doi.org/10.1016/j.anpede.2022.05.011>

* Corresponding author.

E-mail address: pablova.correo@gmail.com

(J.P. Velasco-Amador).

References

1. López-Candela E, Roig Álvaro J, López-Candel J, Fernández Dozagarat S, Soler J, Hernández Bermejo JP, et al. Quistes del rafe medio perineal en la infancia. *An Pediatr.* 2000;52:395-7.
2. Hernández Peláez L, Pontón Martino B, Alonso Losada D, Ordoñez Álvarez FÁ. Quistes del rafe medio: una entidad poco habitual. *Piel.* 2021;36:129.
3. Navalón-Monllor V, Ordoño-Saiz MV, Ordoño-Domínguez F, Sabater-Marco V, Pallás-Costa Y, Navalón-Verdejo P. Quistes de rafe medio en el varón. Presentación de nuestra experiencia y revisión de la literatura. *Actas Urol Esp.* 2017;41:205-9.