

## IMAGES IN PAEDIATRICS

## Phlebectasia of the right jugular vein

## Flebectasia de la vena yugular derecha



María José Sánchez Malo<sup>a,\*</sup>, Lorena Miñones Suárez<sup>a</sup>, Rasha Isabel Perez Ajami<sup>a</sup>,  
María Santos González<sup>b</sup>

<sup>a</sup> Servicio de Pediatría, Hospital Reina Sofía, Tudela, Navarra, Spain


<sup>b</sup> Centro de Salud Cintruénigo, Cintruénigo, Navarra, Spain

Available online 8 June 2024

We present the case of a boy aged 4 years in whom a mass appeared in the right side of the neck when he cried, with onset 1 month prior to consultation. The physical examination was normal while the patient was at rest, but an elongated, soft and compressible painless mass did appear in the right side of the neck upon performance of the Valsalva manoeuvre (Appendix A, video 1).

The cervical ultrasound scan showed a prominent right internal jugular vein, larger in diameter compared to the contralateral vein, that dilated with the Valsalva manoeuvre (Fig. 1). The Doppler ultrasound scan confirmed the presence of venous flow and absence of intraluminal thrombi (Fig. 2).

Jugular vein phlebectasia is a rare condition characterised by a painless, soft and compressible cyst-like/fusiform swelling that appears in the neck of children on crying or straining. Seventy-five percent of cases are diagnosed in children, it is more frequent in boys and it most frequently involves the right brachiocephalic vein.<sup>1–3</sup>



**Figure 1** Ultrasound examination: (A) Right internal jugular vein upon performance of the Valsalva manoeuvre, measuring  $8.2 \times 10.3$  mm. (B) Left jugular vein at rest, with a maximum diameter of 5.8 mm.

It is diagnosed by means of ultrasound and Doppler ultrasound, which allow accurate identification of the disease along with verification of normal venous flow and the potential presence of thrombi.<sup>1,2</sup>

Since this is a benign disorder, most authors recommend conservative management, reserving surgical repair for cases with complications or for cosmetic reasons.<sup>3</sup>


## Funding

This research project did not receive specific financial support from funding agencies in the public, private or not-for-profit sectors.

DOI of original article: <https://doi.org/10.1016/j.anpedi.2024.04.003>

\* Corresponding author.

E-mail address: [mjsanchezmalo@gmail.com](mailto:mjsanchezmalo@gmail.com)  
(M.J. Sánchez Malo).



**Figure 2** Colour Doppler ultrasound: evidence of venous flow and absence of intraluminal thrombi.

## References

1. Girón-Vallejo Ó, del Carmen Benítez-Sánchez M, Villamil V, Hernández-Bermejo JP. A rare (and intermittent) cervical mass: phlebectasia of external jugular vein. *Rev Chil pediatría*. 2017;88(5):688–90.
2. Guerrero Fernández J, Guerrero Vázquez J, García Ascaso MT, Olmedo Sanlaureano S, Luengo Casasola JL. Flebectasia de la vena yugular interna. *An Pediatría*. 2005;63(1):86–8.
3. Jianhong L, Xuewu J, Tingze H. Surgical treatment of jugular vein phlebectasia in children. *Am J Surg*. 2006;192(3):286–90.