



ORIGINAL ARTICLE

Breastfeeding duration and nutritional status of infants and toddlers in Spain. LAYDI study (PAPenRed)



Silvia Martín-Ramos^{a,*}, Begoña Domínguez-Aurrecoechea^b,
Manuela Sánchez Echenique^c, Rubén García Pérez^d, Alicia Bonet Garrosa^e,
Gonzalo Solís-Sánchez^f, en nombre de la Red de Investigación en Pediatría de Atención
Primaria (PAPenRed)¹

^a Servicio de Neonatología del Hospital Río Hortega, Valladolid, Spain

^b Equipo coordinador PAPenRed, Instituto de investigación sanitaria del Principado de Asturias (ISPA), Oviedo, Asturias, Spain

^c Servicio Navarro de Salud Osasunbidea, Servicio de Apoyo a la Gestión Clínica y continuidad asistencial, Pamplona, Spain

^d Txurdinagako Osasun Zentroa, Bilbao, Vizcaya, Spain

^e Centro de Salud Manantiales, Guadalajara, Spain

^f Hospital Universitario Central de Asturias, Instituto de Investigación Sanitaria del Principado de Asturias (ISPA), Universidad de Oviedo, Oviedo, Asturias, Spain

Received 11 May 2024; accepted 12 June 2024

Available online 6 September 2024

KEYWORDS

Breastfeeding;
Growth;
Obesity;
Overweight;
Nutrition

Abstract

Introduction: Breastfeeding appears to protect the onset of obesity in infants. The aim is to know whether breastfeeding duration is associated with the risk of obesity in infants and toddlers aged 12 and 24 months.

Material and methods: Prospective longitudinal study in a cohort of children born in Spain between April 2017 and March 2018 (LAYDI study) in the paediatric primary care system conducted in the framework of the PAPenRed research network. Analysis of breastfeeding duration (group 1: fewer than 6 months; group: more than 6 months) and its association with anthropometric variables.

Results: A total of 1495 patients attended the 12 months preventive child health visit and 1073 patients the 24 months visit. We found a statistically significant association between breastfeeding duration and weight-for-age, BMI-for-age and weight-for-length/height at 12 and 24 months; breastfeeding duration of less than 6 months was significantly associated with overweight and obesity (based on BMI-for-age and weight-for-length/height) at ages 12 and 24 months. Maternal pre-pregnancy BMI acted as a modifier on the association between breastfeeding duration and overweight and obesity (based on BMI-for-age).

DOI of original article: <https://doi.org/10.1016/j.anpedi.2024.06.003>

* Corresponding author.

E-mail address: smartinram@saludcastillayleon.es (S. Martín-Ramos).

¹ Appendix A of the supplemental material lists the paediatricians who collaborated in the LAYDI study.

Conclusions: A breastfeeding duration of less than 6 months is associated with a higher percentage of overweight and obesity at ages 12 and 24 months, although maternal pre-pregnancy BMI modifies this relationship at 24 months.

© 2024 Asociación Española de Pediatría. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Lactancia materna;
Crecimiento;
Obesidad;
Nutrición

Duración de la lactancia materna y condición nutricional de los lactantes en España. Estudio LAYDI (PAPenRed)

Resumen

Introducción: La lactancia materna parece proteger la aparición de la obesidad en los lactantes. El objetivo es conocer si la duración de la lactancia materna se relaciona con el riesgo de obesidad en lactantes de 12 y 24 meses de edad.

Material y métodos: Estudio longitudinal prospectivo de una cohorte de nacidos en España entre abril de 2017 y marzo de 2018 (estudio LAYDI) en pediatría de atención primaria, dentro de la red de investigación PAPenRed. Análisis de duración de lactancia materna (Grupo 1: menos de 6 meses, Grupo: más de 6 meses) y su relación con parámetros somatométricos.

Resultados: Acudieron a la visita de los 12 meses un total de 1495 pacientes, y a la de los 24 meses 1073 pacientes. Se observa una relación estadísticamente significativa entre la duración de la lactancia materna y el peso para la edad, el IMC para la edad y el peso para longitud/talla a los 12 y 24 meses, la duración de la lactancia materna menor de 6 meses se relaciona con el sobrepeso y obesidad (según los índices IMC para la edad y peso para la longitud/talla) de forma estadísticamente significativa a los 12 y 24 meses de edad. El IMC pregestacional materno es una variable modificadora de efecto sobre la relación entre la duración de la lactancia materna y el sobrepeso y obesidad (por IMC para la edad).

Conclusiones: La duración menor de 6 meses de la lactancia materna se asocia a un mayor porcentaje de sobrepeso y obesidad a los 12 y 24 meses, aunque el IMC materno pregestacional modifica esta relación a los 24 meses.

© 2024 Asociación Española de Pediatría. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Early postnatal nutrition is one of the strongest determinants of growth during infancy. Therefore, the role of breastfeeding (BF) in short- and medium-term infant growth and its impact on subsequent development has been assessed in different studies that have yielded results that were not always consistent.^{1–4}

It is known that BF offers benefits in the short, medium and long term as well as in low- to middle-income and high-income developed countries, decreasing overall morbidity and mortality and promoting healthy physical and mental development during childhood and possibly through adulthood.^{5–7}

One of the most intriguing hypothesis in support of BF, especially in developed countries, is that this feeding modality in the early months of life may protect against obesity during childhood and metabolic syndrome in adulthood.^{8–11} Thus, BF could be associated with a lower weight gain in the early months of life compared to formula feeding, which would have significant positive consequences in the medium and long term.

This hypothesis is founded on the potential effect of BF in shaping the gut microbiota, the association between BF

itself and a balanced diet later in childhood or the potential effect of breast milk on the regulation of appetite.^{1,8,12–15} However, other authors argue that the effect of BF in obesity prevention is fundamentally based on socioeconomic factors that may act as confounders in the association between infant feeding modality and obesity.¹⁶

The Mother Infant Study Cohort (MISC) led to the conclusion that in addition to feeding modality, other factors related to the mother (like the maternal body mass index [BMI] or weight gain during pregnancy) may play a key role in this association and have an important impact on the development of malnutrition due to excessive intake during infancy.^{17–19} However, in a very recent publication, Gayle Shipp et al.²⁰ reported that BF acts as a protector against childhood obesity independently of maternal BMI.

The duration of BF seems to be another key factor in this association, so that prolonged BF, whether exclusive or mixed, is associated with decreased weight gain during infancy and childhood, with a stronger effect in the case of exclusive BF.^{4,11,21}

However, the evidence is not that clear. For instance, the impact of BF promotion in pursuit of changes in growth rates and for prevention of obesity in the medium term does not seem consistent and the data on the subject

Table 1 Description of maternal and infant characteristics in the overall cohort and in the 2 groups under study.

	Overall cohort (n = 1946)	Group 1 BF <6 months (n = 997)	Group 2 BF ≥6 months (n = 949)	P (<6 months vs. >6 months)
Female infant	49.9%	50.8%	49.3%	.52 ^a
Maternal BMI, pregestational	24.28 ± 4.78	24.7 ± 5.06	23.8 ± 4.43	<.001 ^b
Pregestational maternal BMI >25	31%	35.5%	26.2%	<.001 ^a
Weight gain during pregnancy (kg)	12.23 ± 5.04	12.35 ± 5.33	12.10 ± 4.72	.26 ^b
Pregnancy weight gain >15 kg	28.6%	29.4%	27.7%	.38 ^a
Gestational age	39.3 ± 1.13	39.3 ± 1.13	39.4 ± 1.13	.06 ^b
Birth weight (kg)	3.30 ± 0.39	3.30 ± 0.39	3.31 ± 0.39	.54 ^b
Residential setting				
Urban	73.7%	71.4%	76.2%	.01 ^a
Rural	21.7%	24.3%	19%	
Mixed	4.6%	4.3%	4.8%	
Type of delivery				
Uncomplicated	68.5%	67.3%	70.4%	.31 ^a
Instrumental	12.8%	13.7%	12%	
Caesarean section	18.3%	19%	17.7%	

BF, breastfeeding; BMI, body mass index.

^a χ^2 test.

^b Student *t* test.

are contradictory.² Furthermore, some authors go as far as questioning whether there is a direct relationship between body composition in infants, whatever the feeding modality, and future health outcomes in the medium and long term.²²

Given the general notion that BF may prevent the development of metabolic complications like obesity in the medium and long term, our primary objective was to analyse the association between the duration of BF and the nutritional status of infants and toddlers in the first 2 years of life as well as the association between the duration of BF and other maternal characteristics (BMI, weight gain during pregnancy).

Material and methods

Prospective longitudinal study in a cohort of children born between April 2017 and March 2018 who attended primary care (PC) paediatric visits in clinics that are part of the Paediatric Primary Care Surveillance Network of Spain (PAPenRED), known as Lactancia y Desarrollo Infantil (Breastfeeding and Infant Development, LAYDI). The PAPenRED is a sentinel network of paediatricians devoted to epidemiological surveillance and research in which PC providers participate on a voluntary basis, while the LAYDI study was designed to understand the impact of socioeconomic, cultural and individual factors on BF initiation and maintenance.

The design and characteristic of the LAYDI study have been described in detail in a previous article,²³ with two-stage stratified sampling throughout the Spanish territory. Each paediatrician selected 12 newborn infants to achieve a total sample size of 1500 infants (out of an estimated finite population of 426 303 newborns in Spain based on birth data for 2015, with an error margin of 2.5% and for a confidence level of 95%).

The inclusion criterion for the LAYDI study was visiting the PC paediatrician during the study period, and for the present study, it was having attended the first and second scheduled well-child visits. We excluded infants born preterm before 37 weeks of gestation, with low birth weight (<2400 g in boys and <2100 g in girls), product of a multiple gestation, with a stay in the maternity or neonatal ward longer than 5 days, with severe disease or malformation, whose mother had moderate to severe health problems during pregnancy or in the postpartum period or whose mothers had difficulty understanding oral or written Spanish.

The LAYDI study analysed variables concerning the pregnancy, childbirth, the neonatal period and social, economic and biological factors. The data were collected at the PC paediatrics clinic in the first routine visit (before 15 days post birth) and successive routine visits at 1, 2, 4, 6, 12, 18 and 24 months. The collaborating paediatricians and nurses collected the data. All anthropometric measurements were made with the certified and calibrated equipment of the respective PC centres.

Among the data collected in the LAYDI study, for this study we retrieved the anthropometric measurements obtained at 12 and 24 months (length in cm and weight in g) and used them to calculate different anthropometric indices, in addition to the infant feeding modality (exclusive breastfeeding [EBF], mixed feeding [MF] and/or formula feeding [FF]). Based on the feeding modality, we defined 2 groups:

- none- Group 1: any feeding modality, as long as BF duration was less than 6 months (including FF, MF and EBF).
- none- Group 2: BF lasting 6 months or longer (including EBF and MF).

Based on the anthropometric measurements, we classified male and female infants by nutritional status based on the World Health Organization (WHO) child growth standards for children under 5 years (WHO 2006).²⁴ We calculated *z*

Table 2 Nutritional status classification of participants based on WHO growth standards at 12 and 24 months.

	12 months	24 months
Weight-for-age z (%)		
Total (n)	1495	1073
≤ -3	1 (0.1%)	1 (0.1%)
-2.99 to -2	16 (1.1%)	9 (0.8%)
-1.99 to -1	148 (9.9%)	109 (10.2%)
-0.99 to +0.99	1077 (72%)	750 (69.9%)
+1 to +1.99	213 (14.2%)	167 (15.6%)
+2 to +2.99	34 (2.3%)	32 (3%)
≥ 3	6 (0.4%)	5 (0.5%)
BMI-for-age z (%)		
Total (n)	1492	1073
≤ -3	2 (0.1%)	4 (0.4%)
-2.99 to -2	13 (0.9%)	18 (1.7%)
-1.99 to -1	149 (10%)	125 (11.6%)
-0.99 to +0.99	1020 (68.4%)	711 (66.3%)
+1 to +1.99	258 (17.3%)	160 (14.9%)
+2 to +2.99	42 (2.8%)	40 (3.7%)
≥ 3	8 (0.5%)	15 (1.4%)
Weight-for-length/height z (%)		
Total (n)	1493	1073
≤ -3	1 (0.1%)	3 (0.3)
-2.99 to -2	13 (0.9%)	11 (1%)
-1.99 to -1	146 (9.8%)	130 (12.1%)
-0.99 to +0.99	1042 (69.8%)	727 (67.8%)
+1 to +1.99	237 (15.9%)	158 (14.7%)
+2 to +2.99	46 (3.1%)	30 (2.8%)
≥ 3	8 (0.5%)	14 (1.3%)
Length/height-for-age z (%)		
Total (n)	1490	1073
≤ -3	9 (0.6%)	8 (0.7%)
-2.99 to -2	40 (2.7%)	23 (2.1%)
-1.99 to -1	197 (13.2%)	132 (12.3%)
-0.99 to +0.99	1017 (68.2%)	712 (66.4%)
+1 to +1.99	192 (12.9%)	167 (15.5%)
+2 to +2.99	35 (2.3%)	28 (2.6%)
≥ 3	2 (0.1%)	4 (0.4%)

BMI, body mass index; WHO, world health organization; z, z score.

scores for the following indicators: weight-for-age, weight for length, length-for-age and BMI-for-age. We grouped z scores based on the following thresholds: less than -3; from -2.99 to -2; from -1.99 to +1.99; from +2 to +2.99, and +3 or greater.

We also classified infants based on their z scores applying the thresholds proposed by the WHO:

- none- Wasting: weight for length/height z score < -2
- none- Stunting: length/height z score < -2.
- none- Underweight: weight-for-age z score less than -2 SD.
- none- Overweight and obesity based on BMI: BMI z score > + 2.
- none- Overweight and obesity based on weight for length/height: weight for length/height z score > +2.

Table 3 Trends in the z scores of nutritional status indicators at ages 12 and 24 months.

	12 months	24 months	P
Weight-for-age z (%)			
Number of cases analysed	1495	1073	
Mean (95% CI)	0.12 (0.07 to 0.17)	0.17 (0.11 to 0.23)	.008
Median (IQR)	0.10 (1.24)	0.22 (1.30)	
SD	0.95	0.96	
BMI z (%)			
Number of cases analysed	1492	1073	
Mean (95% CI)	0.20 (0.15 to 0.25)	0.14 (0.07 to 0.21)	.020
Median (IQR)	0.19 (1.31)	0.14 (1.40)	
SD	0.99	1.10	
Weight-for-length/height z (%)			
Number of cases analysed	1493	1073	
Mean (95% CI)	0.20 (0.15 to 0.25)	0.12 (0.05 to 0.18)	.001
Median (IQR)	0.18 (1.31)	0.10 (1.37)	
SD	0.97	1.05	
Length/height-for-age z (%)			
Number of cases analysed	1490	1073	
Mean (95% CI)	-0.04 (-0.10 to 0.01)	0.10 (0.03 to 0.16)	<.001
Median (IQR)	-0.03 (1.41)	0.12 (1.41)	
SD	1.09	1.06	

Comparison of dependent means. BMI, body mass index; CI, confidence interval; IQR, interquartile range; z, z score.

Lastly, we assessed and quantified a “tendency toward excess weight” category encompassing z scores ranging from +1 to +1.99 for each of the anthropometric parameters.

We have expressed qualitative variables as percentages and quantitative variables as mean and standard deviation (SD). We used the Student *t* test to compare normally distributed continuous variables. For categorical variables, we used the χ^2 test (χ^2 for trend) and for the stratified analysis the Mantel-Haenszel χ^2 test. All the analysis were performed with the statistics package SPSS version 21 (IBM SPSS Statistics) and considering *p* values of less than 0.05 statistically significant.

Results

Description of results in the overall cohort

The sample included 1946 infants. At 1 month of life, 84.9% were breastfed (EBF, 62.3% and MF, 22.6%), a proportion that had decreased to 71.7% at 4 months (EBF, 52.7% and MF, 19%) and to 61.7% at 6 months (EBF, 35.2% and MF, 26.5%). At 12 months, BF continued in 40.1% and at 24 months in 22.3%.

Table 4 Impact at age 12 months of BF duration of less or more than 6 months on the different anthropometric parameters based on z score thresholds.

	Group 1 <6 months	Group 2 >6 months	P ^a
Weight-for-age z (%)			
Total (n)	661	834	<.001
≤ -3	1 (0.2%)	0 (0%)	
-2.99 to -2	5 (0.8%)	11 (1.3%)	
-1.99 to -1	58 (8.8%)	90 (10.8%)	
-0.99 to +0.99	441 (66.7%)	636 (76.3%)	
+1 to +1.99	133 (20.1%)	80 (9.6%)	
+2 to +2.99	21 (3.2%)	13 (1.6%)	
≥ 3	2 (0.3%)	4 (0.5%)	
BMI z (%)			
Total (n)	659	833	<.001
≤ -3	1 (0.2%)	1 (0.1%)	
-2.99 to -2	8 (1.2%)	5 (0.6%)	
-1.99 to -1	52 (7.9%)	97 (11.6%)	
-0.99 to +0.99	428 (64.9%)	592 (71.1%)	
+1 to +1.99	135 (20.5%)	123 (14.8%)	
+2 to +2.99	30 (4.6%)	12 (1.4%)	
≥ 3	5 (0.8%)	3 (0.4%)	
Weight-for-length/height z (%)			
Total (n)	660	833	<.001
≤ -3	1 (0.2%)	0 (0%)	
-2.99 to -2	3 (0.5%)	10 (1.2%)	
-1.99 to -1	55 (8.3%)	91 (10.9%)	
-0.99 to +0.99	430 (65.2%)	612 (73.5%)	
+1 to +1.99	137 (20.8%)	100 (12%)	
+2 to +2.99	29 (4.4%)	17 (2%)	
≥ 3	5 (0.8%)	3 (0.4%)	
Length/height-for-age z (%)			
Total (n)	659	833	.007
≤ -3	5 (0.8%)	4 (0.5%)	
-2.99 to -2	11 (1.7%)	29 (3.5%)	
-1.99 to -1	81 (12.3%)	116 (13.9%)	
-0.99 to +0.99	442 (67.1%)	575 (69%)	
+1 to +1.99	101 (15.3%)	91 (10.9%)	
+2 to +2.99	18 (2.7%)	17 (2%)	
≥ 3	1 (0.2%)	1 (0.1%)	

BF, breastfeeding; CI, confidence interval; z, z score.

^a χ^2 test.

Table 1 presents the general characteristics of the sample, overall and comparing each of the two groups (groups 1 and 2). We found statistically significant differences in the pregestational maternal BMI (greater in group 1) and a significant association with the type of setting (urban/rural/mixed: higher proportion of group 2 in urban settings).

Of the total initial sample, 1495 patients attended the routine visit at age 1 year and 1073 the visit at age 2 years. Of those who attended the visit at 1 year, 44.3% had been breastfed less than 6 months (group 1) and 55.7% 6 or more months (group 2). Of those who attended the visit at 2 years, 44.9% were in group 1 and 55.1% in group 2.

Table 2 provides detailed data on the distribution of the 4 parameters under study (weight-for-age, BMI-for-age, weight for length/height and length/height-for-age) at 12

Table 5 Impact at age 24 months of BF duration of less or more than 6 months on the different anthropometric parameters based on z score thresholds.

	Group 1 <6 months	Group 2 >6 months	P ^a
Weight-for-age z			
Total (n)	481	592	
≤ -3	0 (0%)	1 (0.2%)	<.001
-2.99 to -2	4 (0.8%)	5 (0.8%)	
-1.99 to -1	41 (8.5%)	68 (11.5%)	
-0.99 to +0.99	314 (65.3%)	436 (73.6%)	
+1 to +1.99	97 (20.2%)	70 (11.8%)	
+2 to +2.99	23 (4.8%)	9 (1.5%)	
≥ 3	2 (0.4%)	3 (0.5%)	
BMI z (%)			
Total (n)	481	529	
≤ -3	2 (0.4%)	2 (0.3%)	<.001
-2.99 to -2	5 (1.0%)	13 (2.2%)	
-1.99 to -1	42 (8.7%)	83 (14%)	
-0.99 to +0.99	311 (64.7%)	400 (67.6%)	
+1 to +1.99	85 (17.7%)	75 (12.7%)	
+2 to +2.99	27 (5.6%)	13 (2.2%)	
≥ 3	9 (1.9%)	6 (1.0%)	
Weight-for-length/height z (%)			
Total (n)	481	593	
≤ -3	2 (0.4%)	1 (0.2%)	<.001
-2.99 to -2	3 (0.6%)	8 (1.4%)	
-1.99 to -1	47 (9.8%)	83 (14%)	
-0.99 to +0.99	314 (65.3%)	413 (69.8%)	
+1 to +1.99	87 (18.1%)	71 (12%)	
+2 to +2.99	20 (4.2%)	10 (1.7%)	
≥ 3	8 (1.7%)	6 (1%)	
Length/height-for-age (%)			
Total (n)	481	592	.129
≤ -3	5 (1%)	3 (0.5%)	
-2.99 to -2	9 (1.9%)	14 (2.4%)	
-1.99 to -1	56 (11.6%)	76 (12.8%)	
-0.99 to +0.99	311 (64.7%)	401 (67.7%)	
+1 to +1.99	82 (17%)	84 (14.2%)	
+2 to +2.99	14 (2.9%)	14 (2.4%)	
≥ 3	4 (0.8%)	0 (0%)	

BF, breastfeeding; BMI, body mass index; z, z score.

^a χ^2 test.

and 24 months, and **Table 3** the trends in the z scores between ages 12 and 24 months.

When it came to the low nutritional status categories on the WHO classification, at 12 months wasting (weight for length/height z score < -2) in 1% of the cohort, stunting (length/height z score < -2) in 3.2% and underweight (weight-for-age z score < -2) in 1.2%. At 24 months, these percentages were 1.3%, 2.8% and 0.9%, respectively.

As for the excess weight status categories (WHO), at 12 months, excess weight (overweight or obesity) defined based on the BMI-for-age (z score > +2) was found in the 3.3% of the cohort, and excess weight defined based on weight for length/height (z score > +2) in 3.6%. At 24 months, these percentages were 5.1% and 4.1%, respectively.




Figure 1 Mean values of the weight-for-age, BMI-for-age, weight for length/height and length/height z scores at 12 and 24 months in group 1 and group 2.

When it came to the “tendency toward excess weight”, at 12 months the proportion was 14.2% based on the weight-for-age, 17.3% based on the BMI-for-age, 15.9% based on the weight for length/height and 12.9% based on the length/height-for-age. At 24 months these percentages were 15.6%, 14.9%, 14.7% and 15.5%, respectively.

Association between total BF duration (less and greater than 6 months) and anthropometric values at 12 and 24 months.

At 12 months there was a statistically significant association between the 4 anthropometric parameters under study

and the total duration of BF (Table 4); with a tendency toward higher values in group 1 compared to group 2.

At 24 months there was a statistically significant association between the total duration of BF and the weight-for-age, the BMI-for-age and weight for length/height, but this association was not observed for the length/height-for-age (Table 5). The direction of the trend was the same as at 12 months: greater values in group 1 vs. group 2. Fig. 1 compares the distributions of the 4 analysed anthropometric parameters at 12 and at 24 months based on the duration of BF.

Table 6 Association between WHO categories for abnormal nutritional status (low or excess weight status) and BF duration less or greater than 6 months at ages 12 and 24 months.

	Group 1 (BF duration <6 months)	Group 2 (BF duration >6 months)	<i>P</i>
At 12 months			
Wasting (weight for length/height $z < -2$)	0.6%	1.2%	.237
Stunting (length/height $z < -2$)	2.4%	4.0%	.099
Underweight (weight-for-age $z < -2$)	0.9%	1.3%	.456
Overweight and obesity based on BMI (BMI $z > +2$)	5.3%	1.8%	<.001
Overweight and obesity based on weight for length/height (weight for length/height $z > +2$)	5.2%	2.4%	.005
At 24 months			
Wasting (weight for length/height $z < -2$)	1.0%	1.5%	.490
Stunting (length/height $z < -2$)	2.9%	2.9%	.970
Underweight (weight-for-age $z < -2$)	0.8%	1.0%	.758
Overweight and obesity based on BMI (BMI $z > +2$)	7.5%	3.2%	.002
Overweight and obesity based on weight for length/height (weight for length/height $z > +2$)	5.8%	2.7%	.010

BF, breastfeeding; BMI, body mass index; WHO, World Health Organization; z , z score.

Table 7 Tendency toward excess weight (z score between +1 and +1.99) at ages 12 and 24 months based on breastfeeding duration.

	BF <6 months	BF >6 months	<i>P</i> ^a
At 12 months			
Based on weight-for-age	20.1%	9.6%	<.001
Based on BMI	20.5%	14.8%	.004
Based on weight-for-height	20.8%	12%	<.001
Based on height for age	15.3%	10.9%	.012
At 24 months			
Based on weight-for-age	20.2%	11.8%	<.001
Based on BMI	17.7%	12.7%	.022
Based on weight-for-height	18.1%	12.0%	.005
Based on height for age	17%	14.2%	.198

BF, breastfeeding; BMI, body mass index.

^a Pearson χ^2 test.

Table 6 reflects the impact of BF duration on the WHO nutritional status classification, showing a statistically significant association at both 12 and 24 months between duration of BF of less than 6 months and a higher percentage of excess weight (based on BMI-for-age and weight for length/height).

Table 7 shows the statistically significant association between the duration of BF and the “tendency toward excess weight” at 12 and at 24 months, with a higher percentage in group 1 compared to group 2.

Influence of pregestational maternal weight status on growth

In the study cohort, we found differences in the maternal pregestational BMI between groups 1 and 2 (Table 1), which led us to analyse its impact on nutritional status as a confounder or moderator in the association with the duration of BF.

To assess this effect, we performed a stratified analysis for the association of the duration of BF (<6 months vs. > 6 months) with the classification of excess weight (overweight or obesity) based on the BMI-for-age (z score $\geq +2$ DS), stratifying the maternal pregestational BMI as greater or lesser than 25. We found that the association continued to be significant for a maternal pregestational BMI of less than 25 (OR, 2.9; 95% CI, 1.3–6.6; $P = .006$), but not for a BMI greater than 25 (OR, 1.8; 95% CI, 0.8–4.2; $P = .1439$ [Mantel–Haenszel χ^2 0.005]); thus, the maternal pregestational BMI moderated this association and acted as an interaction variable.

Discussion

The official recommendations and all scientific associations in regard to BF are quite clear, in spite of which current values do not reach the recommended targets.²⁵ The many consequences of poor adherence to BF recommendations, especially in Western societies, seem to include future obesity and metabolic syndrome. Early discontinuation or absence of BF could be related to an increased weight gain in infants in the first months of life, with an impact on childhood and, consequently, on adult life.^{6,26–28}

Excess weight during childhood is a growing health problem in every country, but the study conducted by Rito et al.²⁹ in 22 countries found the highest prevalence in obesity in Spain at 17.7%. There is no question that this problem and its impact on public health are quite serious.

Our study demonstrated a clear association between a duration of BF (in any form) of less than 6 months and a tendency toward greater weight gain at 12 and 24 months based on different anthropometric parameters.

Overall, the 4 parameters under study found a population distribution that clearly fitted the normal distribution. The cohort consisted of healthy boys and girls fed and cared for in a Western society with adequate health care services. The prevalence of the low weight categories in the WHO classification was very low, but the prevalence for excess weight categories was slightly greater.

The longitudinal follow-up of a cohort of this kind allows assessment of the potential impact of infant feeding modal-

ities and their duration on anthropometric values. Thus, we divided the cohort in 2 groups based on the duration of BF (less or more than 6 months), evincing a clear trend: a duration of BF of less than 6 months was associated with increased growth values based on different parameters at both 12 and 24 months. And this trend was objectively demonstrated in the analysis of the percentages categorised as overweight or obesity based on the WHO classification, in addition to the analysis of the separate category of “tendency toward excess weight” that included z scores ranging from +1 to +1.99.

As we commented in the introduction, there are already numerous international studies showing an association between the absence of early discontinuation of BF and excess weight in the first years of life.^{30–32} The study by Ma et al. confirmed that BF can protect against obesity and high body fat²⁹ in children aged 9–11 years in 12 countries and the meta-analysis carried out by the authors of 26 publications adding up to more than 300 000 participants concluded that BF is inversely associated with the risk of early obesity in children aged 2–6 years, with evidence of a dose-response effect between the duration of BF and the reduction in the risk of obesity in early childhood.³² Our study, one of the first conducted on the subject in Spain, corroborates these findings, analysing the routine visits at 1 and 2 years of age and comparing the 4 classic anthropometric parameters in 2 groups defined based on BF duration (less or more than 6 months).

Our study showed that the pregestational maternal weight status (assessed through the BMI) moderates the interaction between BF duration and excess weight (overweight/obesity) based on the BMI at 24 months, so that the maternal pregestational BMI is also variable to consider in this association, as noted by other authors.^{18,20,33}

Conclusion

In our nationwide multicentre prospective cohort study, conducted in the primary care paediatrics setting, a duration of BF of less than 6 months was associated with a greater percentage of excess weight (overweight or obesity) at 12 and 24 months, although there were other factors, such as maternal obesity, that could have interacted with this association.

Ethical considerations

The PAPenRED project was approved by the Ethics and Scientific Research Committee of Aragón (Favourable ruling, minute no 19/2013; C.P.-C.I. P113/00154) and the study presented here was approved by the Research Ethics Committee of the Principality of Asturias (study 213/16). It also received a favourable ruling from the Central Research Committee of the Department of Health of Madrid (February 23, 2017). The mother provided signed informed consent in the first medical visit performed at 15 days post birth. Each child was assigned a code to ensure anonymization.

Funding

The LAYDI study was awarded one of the research grants of the Asociación Española de Pediatría de Atención Primaria-Fundación Pediatría and Salud (AEPap-FPS) in 2017.

Conflicts of interest

The authors have no conflicts of interest to declare.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.anpede.2024.08.003>.

References

- Horta BL, Rollins N, Dias MS, Garcez V, Pérez-Escamilla R. Systematic review and meta-analysis of breastfeeding and later overweight or obesity expands on previous study for World Health Organization. *Acta Paediatr.* 2023;112(1):34–41, <http://dx.doi.org/10.1111/apa.16460>. Epub 2022 Jul 1. PMID: 35727183.
- Giugliani ER, Horta BL, Loret de Mola C, Lisboa BO, Victora CG. Effect of breastfeeding promotion interventions on child growth: a systematic review and meta-analysis. *Acta Paediatr.* 2015;104(467):20–9, <http://dx.doi.org/10.1111/apa.13160>.
- Dewey K, Bazzano L, Davis T, Donovan S, Taveras E, Kleinman R, et al. The Duration, Frequency, and Volume of Exclusive Human Milk and/or Infant Formula Consumption and Overweight and Obesity: A Systematic Review [Internet]. Alexandria (VA): USDA Nutrition Evidence Systematic Review; 2020.
- Qiao J, Dai LJ, Zhang Q, Ouyang YQ. A meta-analysis of the association between breastfeeding and early childhood obesity. *J PediatrNurs.* 2020;53:57–66, <http://dx.doi.org/10.1016/j.pedn.2020.04.024>.
- Victora CG, Bahl R, Barros AJ, França GV, Horton S, Krasevec J, et al., Lancet Breastfeeding Series Group. Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. *Lancet.* 2016;387(10017):475–90, [http://dx.doi.org/10.1016/S0140-6736\(15\)01024-7](http://dx.doi.org/10.1016/S0140-6736(15)01024-7).
- Horta BL, Loret de Mola C, Victora CG. Breastfeeding and intelligence: a systematic review and meta-analysis. *Acta Paediatr.* 2015;104:14–9.
- Wallenborn JT, Levine GA, Carreira Dos Santos A, Grisi S, Brentani A, Fink G. Breastfeeding, physical growth, and cognitive development. *Pediatrics.* 2021;147(5):e2020008029, <http://dx.doi.org/10.1542/peds.2020-008029>.
- Horta BL, Loret de Mola C, Victora CG. Long-term consequences of breastfeeding on cholesterol, obesity, systolic blood pressure and type 2 diabetes: a systematic review and meta-analysis. *Acta Paediatr.* 2015;104(467):30–7, <http://dx.doi.org/10.1111/apa.13133>.
- Mazur D, Satora M, Rekowska AK, Kabała Z, Łomża A, Kimber-Trojnar Z, Leszczyńska-Gorzela B. Influence of breastfeeding on the state of meta-inflammation in obesity-A narrative review. *Curr Issues Mol Biol.* 2023;45(11):9003–18, <http://dx.doi.org/10.3390/cimb45110565>.
- Zheng M, D'Souza NJ, Atkins L, Ghobadi S, Laws R, Szymlek-Gay EA, et al. Breastfeeding and the longitudinal changes of body mass index in childhood and adulthood: a systematic review. *AdvNutr.* 2024;15(1):100152, <http://dx.doi.org/10.1016/j.advnut.2023.100152>.
- Aguilar Cordero MJ, Sánchez López AM, Madrid Baños N, Mur Villar N, Expósito Ruiz M, Hermoso Rodríguez E. Lactancia materna como prevención del sobrepeso y la obesidad en el niño y el adolescente; revisión sistemática. *NutrHosp.* 2014;31(2):606–20, <http://dx.doi.org/10.3305/nh.2015.31.2.8458>.
- Selma-Royo M, Calvo Lerma J, Cortés-Macías E, Collado MC. Human milk microbiome: from actual knowledge to future perspective. *Semin Perinatol.* 2021;45(6):151450, <http://dx.doi.org/10.1016/j.semperi.2021.151450>.
- Juan Castell MF, Peraita-Costa I, Soriano JM, Llopis-Morales A, Morales-Suarez-Varela M. A review of the relationship between the appetite-regulating hormone leptin present in human milk and infant growth. *Breastfeed Med.* 2022;17(2):98–111, <http://dx.doi.org/10.1089/bfm.2021.0149>.
- Taveras EM. Association of breastfeeding with maternal control of infant feeding at age 1 year. *Pediatrics.* 2004;114(5):e577–83.
- Cortés-Macías E, Selma-Royo M, Rio-Aige K, Bäuerl C, Rodríguez-Lagunas MJ, Martínez-Costa C, et al. Distinct breast milk microbiota, cytokine, and adipokine profiles are associated with infant growth at 12 months: an in vitro host-microbe interaction mechanistic approach. *Food Funct.* 2023;14(1):148–59, <http://dx.doi.org/10.1039/d2fo02060b>.
- Brion MJA, Lawlor DA, Matijasevich A, Horta B, Anselmi L, Araujo CL, et al. What are the causal effects of breastfeeding on IQ, obesity and blood pressure? Evidence from comparing high-income with middle-income cohorts. *Int J Epidemiol.* 2011;40(3):670–80.
- Radwan H, Hashim M, Shaker Obaid R, Hasan H, Naja F, Al Ghazal H, et al. The Mother-Infant Study Cohort (MISC): methodology, challenges, and baseline characteristics. *PLoS One.* 2018;13(5):e0198278, <http://dx.doi.org/10.1371/journal.pone.0198278>.
- Radwan H, Naja F, Abbas N, Kharma JA, Obaid RS, Al Ghazal H, et al. Influence of maternal characteristics and infant feeding patterns on infant growth from birth till 18 months: The MISC study. *PediatrObes.* 2024:e13106, <http://dx.doi.org/10.1111/ijpo.13106>. Epub ahead of print.
- Enstad S, Cheema S, Thomas R, Fichorova RN, Martin CR, O'Tierney-Ginn P, et al. The impact of maternal obesity and breast milk inflammation on developmental programming of infant growth. *Eur J Clin Nutr.* 2021;75(1):180–8, <http://dx.doi.org/10.1038/s41430-020-00720-5>.
- Shipp GM, Wosu AC, Knapp EA, Sauder KA, Dabelea D, Perng W, et al., program collaborators for Environmental influences on Child Health Outcomes. Maternal pre-pregnancy BMI, breastfeeding, and child BMI. *Pediatrics.* 2024;153(1):e2023061466, <http://dx.doi.org/10.1542/peds.2023-061466>.
- Patro-Gołąb B, Zalewski BM, Polaczek A, Szajewska H. Duration of breastfeeding and early growth: a systematic review of current evidence. *Breastfeed Med.* 2019;14(4):218–29, <http://dx.doi.org/10.1089/bfm.2018.0187>.
- Amati F, McCann L, Castañeda-Gutiérrez E, Prior E, van Loo-Bouwman CA, Abrahamse-Berkeveld M, et al. Infant fat mass and later child and adolescent health outcomes: a systematic review. *ArchDis Child.* 2024;109(2):125–9, <http://dx.doi.org/10.1136/archdischild-2023-325798>.
- Martin-Ramos S, Domínguez-Aurrecoechea B, García Vera C, Lorente García Mauriño AM, Sánchez Almeida E, Solís-Sánchez G. Lactancia materna en España y factores relacionados con su instauración y mantenimiento: estudio LAYDI (PAPenRed). *Aten Primaria.* 2024;56(1):102772, <http://dx.doi.org/10.1016/j.aprim.2023.102772>.
- WHO. The Multicentre Growth Reference Study Group: WHO Child Growth Standards Length/Height-for-Age, Weight-for-Age, Weight-for-Length, Weight-for-Height and Body Mass Index for-Age: Methods and Development. 2006 Last accessed on 2020 Jun 03 Geneva WHO Available from:

- <https://www.who.int/toolkits/child-growth-standards/standards/weight-for-age>.
25. World Health Organization. Global nutrition targets 2025: breastfeeding policy brief (WHO/NMH/NHD/14.7) [Internet]. WHO/UNICEF, (2014), [consulta: 17 de noviembre de 2022] Available from: <https://www.who.int/publications/i/item/WHO-NMH-NHD-14.7>.
 26. Binns C, Lee M, Low WY. The long-term public health benefits of breastfeeding. *Asia Pac J Public Health*. 2016;28:7.
 27. Wijnhoven TM, van Raaij JM, Spinelli A, Starc G, Hassapidou M, Spiroski I, et al. WHO European Childhood Obesity Surveillance Initiative: body mass index and level of overweight among 6-9-year-old children from school year 2007/2008 to school year 2009/2010. *BMC Public Health*. 2014;14(1):806.
 28. Eidelman AI. Breastfeeding goals: are we fighting the right war? *Breastfeed Med*. 2021;16:929-30.
 29. Rito AI, Buoncristiano M, Spinelli A, Salanave B, Kunešová M, Hejgaard T, et al. Association between Characteristics at birth, breastfeeding and obesity in 22 countries: the WHO European Childhood Obesity Surveillance Initiative - COSI 2015/2017. *Obes Facts*. 2019;12(2):226-43, <http://dx.doi.org/10.1159/000500425>.
 30. Jwa SC, Fujiwara T, Kondo N. Latent protective effects of breastfeeding on late childhood overweight and obesity: a nationwide prospective study. *Obesity (Silver Spring)*. 2014;22(6):1527-37, <http://dx.doi.org/10.1002/oby.20735>.
 31. Marseglia L, Manti S, D'Angelo G, Cuppari C, Salpietro V, Filippelli M, et al. Obesity and breastfeeding: the strength of association. *Women Birth*. 2015;28(2):81-6, <http://dx.doi.org/10.1016/j.wombi.2014.12.007>.
 32. Ma J, Qiao Y, Zhao P, et al. Breastfeeding and childhood obesity: a 12-country study. *Matern Child Nutr*. 2020;16(3):e12984, <http://dx.doi.org/10.1111/mcn.12984>.
 33. Iglesia Altaba I, Larqué E, Mesa MD, Blanco-Carnero JE, Gomez-Llorente C, Rodríguez-Martínez G, et al. Early nutrition and later excess adiposity during childhood: a narrative review. *Horm Res Paediatr*. 2022;95(2):112-9, <http://dx.doi.org/10.1159/000520811>.