

IMAGES IN PAEDIATRICS

Fatal gastric perforation caused by *Chlostridium ventriculi*. A rare presentation of abdominal pain



Perforación gástrica letal secundaria a *Chlostridium ventriculi*. Una causa inusual de dolor abdominal

Felipe Stacey^a, Pablo Moreno Santabarbara^b, Andrea Feu Llauradó^c,
Javier Rodríguez-Fanjul^{a,*}

^a Unidad de Cuidados Intensivos Pediátricos, Servicio de Pediatría, Hospital Germans Trias i Pujol, Badalona, Spain

^b Unidad Endocrina, Metabólica y Bariátrica, Servicio de Cirugía General, Hospital Germans Trias i Pujol, Badalona, Spain

^c Servicio de Patología Patológica, Hospital Germans Trias i Pujol, Badalona, Spain

Received 22 July 2024; accepted 4 October 2024

Available online 27 January 2025

A previously healthy boy aged 13 years was admitted with sudden acute abdominal pain. The family reported weekly nocturnal vomiting episodes over a 6-month period.

At admission, the patient was in shock, with a distended and rigid abdomen, respiratory failure and decreased level of consciousness due to global hypoperfusion secondary to abdominal compartment syndrome. This prompted intubation and initiation of fluid resuscitation and vasoactive drugs. The abdominal radiograph showed massive gastric distension with small and large bowel displacement. An emergency exploratory laparotomy revealed multiple gastric perforations, leading to performance of partial gastrectomy (Fig. 1). The antimicrobial spectrum was broadened, with administration of carbapenems, metronidazole, vancomycin and antifungals. The patient underwent pro-



Figure 1 Macroscopic appearance of gastric specimen from partial gastrectomy. Multiple gastric perforations.

gressive deterioration with multiple organ failure, including acute kidney injury requiring continuous renal replacement and acute liver failure requiring therapeutic plasma exchange.

DOI of original article:

<https://doi.org/10.1016/j.anpedi.2024.503741>

* Corresponding author.

E-mail address: javier.rodriquez.fanjul@gmail.com

(J. Rodríguez-Fanjul).




Figure 2 Histological examination of the surgical gastric specimen. Standard haematoxylin-eosin staining revealed diffuse haemorrhagic areas with extensive ischaemic necrosis of the mucosa and submucosa and the presence of *C ventriculi* bacteria with characteristic microscopic features: spherical basophilic bacteria (cocci), each measuring 1.8 to 3 μm , arranged in tetrads or octets forming cuboid structures.

The histological examination of gastric specimen revealed pathogenic colonization by *Clostridium ventriculi* (Fig. 2). In spite of the care provided, the patient died.

Clostridium ventriculi (also known as *Sarcina ventriculi*) is an anaerobic gram-positive coccus mainly transmitted through ingestion of contaminated soil present in food.¹ The main risk factor is delayed gastric emptying. Intramural gas accumulation from bacterial fermentation byproducts results in emphysematous gastritis, with a mortality of 75%.² Very few paediatric cases have been reported to date,³ and the one we present here is the first with a fatal outcome.

The diagnosis can be made with the observation of the tetrad packet arrangement of *C ventriculi* with haematoxylin-eosin staining (Fig. 2).

References

1. Rizwan M, Kim G, Senseng C. *Sarcina ventriculi*: review of the Literature. *Arch Pathol Lab Med.* 2016;140:1441–5.
2. Singh K. Emphysematous gastritis associated with *Sarcina ventriculi*. *Case Rep Gastroenterol.* 2019;13:207–13.
3. Marcelino L, Valentini Junior D, dos Santos Machado SM, Guilherme Schaefer P, Camara Rivero R, Bersch Osvaldt A. *Sarcina ventriculi* a rare pathogen. *Autops Case Rep.* 2021;11:1–10.