

IMAGES IN PAEDIATRICS

Ileocecal intussusception secondary to heterotopic pancreas**Invaginación ileocecal secundaria a páncreas heterotópico**

Antonio Palacios Prados^{a,*}, Rocío Vizcaíno Pérez^a, Ana López Prieto^b,
Madai Curbelo Rodríguez^a

^a Servicio de Cirugía Pediátrica, Hospital Universitario Virgen Macarena, Sevilla, Spain

^b Servicio de Anatomía Patológica, Hospital Universitario Virgen Macarena, Sevilla, Spain



Figure 1 Ultrasound scan showing ileocecal intussusception.

A boy aged 16 months presented in the emergency department with asthenia, abdominal pain and vomiting of 72 h' duration. The key finding of the physical examination

was the detection on palpation of an abdominal mass in the umbilical region.

The mass was evaluated with an abdominal ultrasound scan (Fig. 1), leading to diagnosis of advanced ileocecal intussusception and possible bowel perforation.

The patient underwent urgent surgery via midline laparotomy, which revealed ileocecal intussusception extending to the hepatic flexure (Fig. 2A).

Reduction of the intussusception made visible a necrotic ileal segment with a round, indurated lesion that acted as

DOI of original article:

<https://doi.org/10.1016/j.anpedi.2025.503805>

* Corresponding author.

E-mail address: antonio.palacios.prados@hotmail.com

(A. Palacios Prados).




Figure 2 Intraoperative images.

Top: ileocecal intussusception. Bottom left: necrotic bowel segment. Bottom right: round indurated lesion compatible with heterotopic pancreas.

the head of intussusception (Figs. 2B and C). The affected bowel was resected, followed by end-to-end anastomosis, and the surgical specimen submitted for pathological examination.

The histological study confirmed the presence of heterotopic pancreas at the level of the ileum (Fig. 3), which had caused the intussusception.

The patient did not experience any postoperative complications.

Heterotopic or ectopic pancreas is defined as the presence of pancreatic tissue outside the normal anatomical location of the pancreas.¹ It is a rare condition in the pediatric population that, while usually asymptomatic, may manifest in the form of pancreatitis, infection, malignancy or intussusception (with heterotopic pancreas accounting for 1%–2% of total cases of intussusception).²




Figure 3 Histological study.

Top left: transmural lesion extending from the mucosa to the serous layer, composed of pancreatic ducts and acini but without islets of Langerhans. Top right: at greater magnification, pancreatic acini could be visualized, composed of polygonal cells without atypia. Bottom: immunohistochemical staining was positive for neuroendocrine markers such as chromogranin (left) and synaptophysin (right).

References

1. Bazán Zender C, Reyes Coloma L, León Cueto JL, Revoredo Palacios G, Stella Castillo JA, Pezo A. Páncreas heterotópico como causa de invaginación intestinal: primer caso reportado en el Perú. *Rev Perú Med Exp Salud Publica*. 2015;32:598–602
2. Galván-Montaño A, Suárez-Roa ML, Estrada-Hernández MR. Invaginación yeyuno-yeyunal secundaria a tejido pancreático ectópico en un lactante de un año de edad. Reporte de caso. *Cir Cir*. 2012;80:546–9. <http://www.scielo.org.pe/scielo.php?script=sci.arttext&pid=S1726-46342015000300027&lng=es>