



ORIGINAL ARTICLE

Prevalence and temporal evolution of asthma symptoms in Spain. Global Asthma Network (GAN) study[☆]



Alberto Bercedo Sanz^{a,b,*}, Antonela Martínez-Torres^{c,d}, Carlos González Díaz^e, Ángel López-Silvarrey Varela^f, Francisco Javier Pellegrini Belinchón^{g,h}, Inés Aguinaga-Ontoso^{i,j}, Luis García-Marcos^{d,k}, Grupo GAN España¹

^a Centro de Salud Dobra, Torrelavega, Servicio Cántabro de Salud, Cantabria, Spain

^b Instituto de Investigación Sanitaria Valdecilla, IDIVAL, Santander. Cantabria, Spain

^c Unidad de Neumología y Alergia Pediátrica y Grupo de Investigación en Enfermería, Hospital Infantil Universitario Virgen de la Arrixaca, Murcia, Spain

^d Instituto Murciano de Investigación Biosanitaria IMIB, ARADyAL Allergy Network, Murcia, Spain

^e Unidad de Alergia Infantil, Hospital Universitario Basurto, Bilbao, Vizcaya, Spain

^f Fundación María José Jove, Servicio Galego de Saúde (SERGAS), La Coruña, Spain

^g Centro de Salud Pizarrales, Salamanca, Spain

^h Departamento de Ciencias Biomédicas y del Diagnóstico, Universidad de Salamanca, Salamanca, Spain

ⁱ Departamento de Ciencias de la Salud, Universidad Pública de Navarra (UPNA), Pamplona, Spain

^j IdiSNA, Instituto de Investigación Sanitaria de Navarra, Pamplona, Spain

^k Unidad de Neumología y Alergia Pediátrica, Hospital Infantil Universitario Virgen de la Arrixaca, Universidad de Murcia, Murcia, Spain

Received 16 August 2021; accepted 20 October 2021

Available online 26 July 2022

KEYWORDS

Asthma;
Adolescent;
Schoolchild;
Prevalence;
Cross-sectional
studies;
Global Asthma
Network

Abstract

Introduction: The temporal evolution of the prevalence of asthma described in the ISAAC (International Study of Asthma and Allergies in Childhood) in 2002 is unknown, or if the geographical or age differences are maintained in Spain.

Objective: To describe the prevalence of asthma symptoms in different Spanish geographic areas and compare it with that of those centers that participated in the ISAAC.

Methods: Cross-sectional study of asthma prevalence, carried out in 2016–2019 with 19,943 adolescents aged 13–14 years and 17,215 schoolchildren aged 6–7 years from 6 Spanish geographical areas (Cartagena, Bilbao, Cantabria, La Coruña, Pamplona and Salamanca). Asthma symptoms were collected using a written questionnaire and video questionnaire according to the Global Asthma Network (GAN) protocol.

[☆] Please cite this article as: Bercedo Sanz A, Martínez-Torres A, González Díaz C, López-Silvarrey Varela A, Pellegrini Belinchón FJ, Aguinaga-Ontoso I, et al. Prevalencia y evolución temporal de síntomas de asma en España. Estudio *Global Asthma Network* (GAN). *An Pediatr (Barc)*. 2022;97:161–171.

* Corresponding author.

E-mail addresses: drbercedo@gmail.com, alberto.bercedo@scsalud.es (A. Bercedo Sanz).

¹ Appendix A lists the members of the GAN Spain group.

Results: The prevalence of recent wheezing (last 12 months) was 15.3% at 13–14 years and 10.4% at 6–7 years, with variations in adolescents, from 19% in Bilbao to 10.2% in Cartagena; and in schoolchildren, from 11.7% in Cartagena to 7% in Pamplona. These prevalences were higher than those of the ISAAC (10.6% in adolescents and 9.9% in schoolchildren). 21.3% of adolescents and 12.4% of schoolchildren reported asthma at some time.

Conclusions: There is a high prevalence of asthmatic symptoms with an increase in adolescents and a stabilization in Spanish schoolchildren with respect to the ISAAC. Geographic variations in asthma prevalence are not so clearly appreciated, but areas with high prevalences maintain high numbers.

© 2021 Asociación Española de Pediatría. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

PALABRAS CLAVE

Asma;
Adolescente;
Escolar;
Prevalencia;
Estudios
transversales;
Global Asthma
Network

Prevalencia y evolución temporal de síntomas de asma en España. Estudio *Global Asthma Network* (GAN)

Resumen

Introducción: Se desconoce la evolución temporal de la prevalencia de asma descrita en el ISAAC (International Study of Asthma and Allergies in Childhood) en 2002 o si las diferencias geográficas o por edades se mantienen en España.

Objetivo: Describir la prevalencia de los síntomas de asma en distintas áreas geográficas españolas y compararla con la de aquellos centros que participaron en el ISAAC.

Métodos: Estudio transversal de prevalencia de asma, realizado en 2016–2019 a 19,943 adolescentes de 13–14 años y 17,215 escolares de 6–7 años de 6 áreas geográficas españolas (Cartagena, Bilbao, Cantabria, La Coruña, Pamplona y Salamanca). Los síntomas de asma se recogieron mediante un cuestionario escrito y videocuestionario según el protocolo Global Asthma Network (GAN).

Resultados: La prevalencia de sibilancias recientes (últimos 12 meses) fue del 15,3% a los 13–14 años y del 10,4% a los 6–7 años, con variaciones en los adolescentes, desde un 19% en Bilbao hasta un 10,2% en Cartagena; y en los escolares, desde un 11,7% en Cartagena hasta un 7% en Pamplona. Estas prevalencias fueron superiores a las del ISAAC (10,6% en adolescentes y 9,9% en los escolares). Un 21,3% de adolescentes y un 12,4% de los escolares refirieron asma alguna vez.

Conclusiones: Existe una alta prevalencia de síntomas asmáticos con un incremento en los adolescentes y una estabilización en los escolares españoles con respecto al ISAAC. No se aprecian tan claramente variaciones geográficas en la prevalencia de asma, pero las áreas que tenían prevalencias elevadas mantienen cifras altas.

© 2021 Asociación Española de Pediatría. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Asthma is a common chronic disease in childhood and adolescence, with a prevalence that varies between geographical regions and age groups.^{1,2} According to the Global Burden of Disease study, asthma is the leading cause of disability, measured in disability-adjusted life years (DALYs), in children aged 5–14 years in developed countries.³

The International Study of Asthma and Allergies in Childhood (ISAAC, 1992–2012) has contributed a vast collection of global epidemiological data on aspects such as prevalence, risk factors and protective factors (<http://isaac.auckland.ac.nz/>), as well as the identification of candidate genes for wheezing and allergy⁴ and different asthma phenotypes.⁵ The first data on the prevalence of asthma in Spain collected in multiple centres with a shared, standardised and validated methodology became available in 1998. The data showed a prevalence of asthma in

the past year of 9.3% in children aged 13–14 years and of 6.2% in children aged 6–7 years.¹

The data obtained in phase III of the ISAAC study (ISAAC-III, 2002–2003) in Spain suggested that the Mediterranean diet may be protective against asthma in school-age children and adolescents,^{6–8} in addition to highlighting the relevance of obesity in this disease.⁹ This phase also established 2 thresholds for the prevalence of asthma, confirming an increase to up to 9.9% in children aged 6–7 years and stabilization at 10.6% in adolescents, with identification of a high-prevalence geographical area corresponding to Green Spain (the ‘‘cornisa cantábrica’’) with a prevalence in adolescents of 12.8% in Bilbao, 15.2% in A Coruña, 15.3% in Asturias and 16.7% in Cantabria.^{10–12} Other relevant findings included variations associated with climate^{13,14} and the considerable impact of vaccination with bacillus Calmette-Guérin (BCG)¹⁵ or urban environmental pollution¹⁶ in the prevalence of asthma in Spain.

At present, we do not know if the observed temporal trends in the prevalence of asthma, with a plateau in adolescents and an increase in younger children, or the geographical differences described in the ISAAC-III have been sustained. Furthermore, while the prevalence of asthma and the associated risk or protective factors are well defined in some instances, these factors may behave differently in different locations and at different times.

In this context, and applying the ISAAC methodology, the Global Asthma Network (GAN) study (www.globalasthmanetwork.org) was launched with the aim of not only updating the information on the prevalence of asthma and the associated factors globally, but also to assess the associated costs and the approaches to its management. At present, the GAN network comprises 55 centres in 20 countries, including 6 in Spain (Murcia, Bilbao, Cantabria, La Coruña, Pamplona and Salamanca).^{17,18}

The objective of our study was to describe the current prevalence of asthma symptoms in GAN centres in Spain and compare it with the prevalence found in the centres that participated in the ISAAC-III.

Methods

We conducted a cross-sectional study applying the methodology described in the Global Asthma Network Manual for Global Surveillance (<http://globalasthmanetwork.org/surveillance/manual/manual.php>). It was based on data collected through standardised and previously validated questionnaires (<http://globalasthmanetwork.org/surveillance/manual/validation.php>) in a general sample, with an estimated sample size of 3000 participants per age group (13/14 years and 6/7 years) required to allow detection of differences in asthma prevalence of 3% or greater with a power of 90% and a level of confidence of 99%.^{17,18}

The field work took place during the school year at different points in the 2016–2019 period, depending on the centre (Table 1). The principal investigator of each participating centre contacted the corresponding Department of Education to obtain authorization to carry out the study in the selected primary and secondary schools, and subsequently communicated with the school principals, teachers and parents to inform them of the nature of the study. The sample included students enrolled in years 2 and 3 of compulsory secondary education (*educación secundaria obligatoria*, ESO) (aged 13/14 years) or years 1 and 2 of primary education (aged 6/7 years) in public or private schools. A schedule of visits to participating schools was established to guide the distribution and submission of self-administered paper- and video-based questionnaires completed by adolescents and the paper-based questionnaires for children aged 6/7 years completed by parents, and to oversee the entire process.

The main instrument in data collection was the validated questionnaire, which includes sections on the symptoms of asthma and other allergic diseases, their severity, use of resources, risk and/or protective factors and medication. The question was translated from English to Spanish and back-translated to English applying the ISAAC methodology to ensure that the original meaning of the items was retained.¹⁹ Adolescents also completed a video questionnaire, translated to Spanish, that includes 5 scenarios in which individuals of different ages experience asthma exacerbations of varying severity and in different contexts. The Spanish version can be downloaded at http://pediatria.imib.es/portal/instituto/pediatria_gan.jsf, and the specific variables under study and their coding at <http://globalasthmanetwork.org/surveillance/manual/coding.php>. The primary variable was the presence of wheezing

in the past year, defined as a ‘yes’ answer to the item asking whether the child or adolescent had experienced wheezing or whistling in the chest in the past 12 months. Severe wheeze was defined as having experienced 4 or more wheezing attacks, or 1 or more awakenings at night due to wheezing per week, or experiencing wheezing severe enough to limit speech, all of it in reference to the past 12 months. A positive history of asthma was defined as a ‘yes’ answer to ‘‘Have you/had this child ever had asthma?’’.

The questionnaires were anonymised and scanned for reading in the coordinating centre in Cartagena (Murcia) with a Fujitsu fi-7700 scanner with optical mark recognition (OMR) software (Remark Office OMR version 10; Gravic Inc; Malvern, PA, USA). We performed a descriptive analysis of the variables followed by a comparative bivariate analysis using the chi square test to estimate possible differences based on sex in participating centres. Results were considered statistically significant if the *P* value was 0.05 or less. The statistical analysis was performed with the software STATA version 15 (College Station, TX, USA).

The GAN study was approved in the national coordinating centre by the Clinical Research Ethics Committee of the Hospital Clínico Universitario Virgen de la Arrixaca in Murcia, and subsequently validated and approved in each participating centre. In adolescents, we obtained informed consent from the parents.

Results

The GAN study in Spain involved the participation of 6 centres with a sample of 19 943 adolescents aged 13–14 years and 17 215 schoolchildren aged 6–7 years. The sample of adolescents was recruited from 184 schools and the sample of schoolchildren aged 6–7 years from 323 schools (Table 1). Participation was higher in the 13/14 years group (overall, 81.3%; range, 58.8%–95%) compared to the 6/7 years group (overall, 62.6%; range, 55.2%–73.7%).

Tables 2 and 3 present the prevalence figures obtained through the paper-based questionnaire in each age group. The overall prevalence of recent wheezing for the total of participating GAN centres was 15.3% in adolescents and 10.4% in schoolchildren. As regards variation, the prevalence in adolescents ranged from 10.2% in Cartagena to 19% in Bilbao. There was less variation in schoolchildren, with a maximum of 11.7% in Cartagena and a minimum of 7% in Pamplona. In the group aged 6/7 years, only Salamanca and Pamplona had a prevalence under 10%. The prevalence of a positive history of asthma (ever had asthma) was also high: 21.3% in adolescents and 12.4% in schoolchildren.

As regards variables associated with asthma severity, we found a prevalence of wheezing impeding speech of 4.3% in adolescents and 2% in schoolchildren. The proportions of patients that reported 4–12 and more than 12 wheezing attacks in the past year were 3.6% and 1.4%, respectively, in adolescents, and 2% and 0.5%, respectively, in schoolchildren. The prevalence of sleep disturbances (awakening due to wheezing 1 or more nights per week in the recent past) was 1.3% in adolescents and 1.5% in schoolchildren.

The analysis of asthma symptoms assessed through the video questionnaire in adolescents (Table 4) yielded a lower prevalence of recent wheezing compared to the paper-based questionnaire in every centre, with an overall prevalence of 12.9%, and differences between the 2 questionnaires ranging from a minimum of 7.5% in Cartagena to a maximum of 17.2% in Bilbao. This was not the case when we compared the prevalence of severe asthma exacerbations obtained with the 2 instruments, as the pattern went in the other direction, with a lower prevalence obtained with the video questionnaire compared to

Table 1 Characteristics of participating GAN centres.

GAN centre	Setting	Participating schools	Eligible students	Sample size	Response rate (% eligible students)
<i>Bilbao</i>					
6–7 years	Urban	52	4902	2707	55.2%
13–14 years		33	3711	3379	91.1%
<i>Cantabria</i>					
6–7 years	Urban	75	5052	2841	56.2%
13–14 years		47	5664	4382	77.3%
<i>Cartagena</i>					
6–7 years	Urban	61	5342	3509	65.7%
13–14 years		26	4657	3437	73.8%
<i>A Coruña</i>					
6–7 years	Urban	48	4796	3407	71.0%
13–14 years		26	3760	3462	92.1%
<i>Pamplona</i>					
6–7 years	Urban	36	4163	2363	56.8%
13–14 years		21	3056	1798	58.8%
<i>Salamanca</i>					
6–7 years	Mixed	51	3242	2388	73.7%
13–14 years		31	3668	3485	95.0%
<i>Total</i>					
6–7 years		323	27 497	17 215	62.6%
13–14 years		184	24 516	19 943	81.3%

the paper-based one (8.5% vs 4.3%). When it came to the sleep disturbances caused by wheezing in the past 12 months, there was no overall difference in prevalence between the 2 instruments (video questionnaire, 6% vs paper questionnaire, 6.3%). The prevalence of cough at night was substantially lower in the video questionnaire compared to the paper-based one (21.9% vs 31%), while there was vary any difference in prevalence of recent wheezing associated with exercise (20.8% in the video questionnaire vs 21.7% in the paper questionnaire).

Tables 5 and 6 compare the prevalence of asthma symptoms in the past year obtained in the ISAAC-III (2002–2003) and the GAN study (2016–2019) in the 2 age groups. In adolescents, the prevalence of wheezing in the past year was higher in female adolescents in all GAN centres, with the opposite trend in schoolchildren, in which the prevalence was higher in boys. In some GAN centres, the observed difference based on sex was greater compared to the ISAAC-III. We also found a significant increase in the GAN study in the prevalence of wheezing in the past year in adolescents (15.3% vs 12.2%) and a stable prevalence in schoolchildren (10.4% vs 10.7%). When it came to severe wheezing, we found a significant increase in prevalence in adolescents (7% vs 5.1%) and no difference in schoolchildren (3.8% vs 3.7%).

Discussion

The GAN study has contributed updated information in the prevalence of asthma symptoms in Spain, confirming that asthma continues to be a common affliction with a prevalence of wheezing in the recent past of 15.3% in adolescents and 10.4% in schoolchildren, reflecting an increase compared to the prevalence in 2002–2003 found in the ISAAC-III (10.6% and 9.9%, respectively).^{10,11} While the prevalence exhibited an increase in adolescents, it appeared to be plateauing in children. While the increase in adolescents may be partly explained by the smaller number of centres that participated in the GAN study and the

absence of some of the centres included in the ISAAC-III (Castellon, Barcelona and Valladolid) where the prevalence of asthma had been low, it is likely that the observed increase was real, as the prevalence in the GAN centre in Pamplona, which also participated in the ISAAC-III, increased from 8% to 16.9%, and the GAN centre in Salamanca, which did not participate in the ISAAC-III, reported a current prevalence of 14.7%, much higher compared to the past prevalence in nearby centres, such as the one in Valladolid (prevalence of 8.2% in the ISAAC-III).

Our findings only partially confirmed the results of the ISAAC-III concerning geographical variations with a high-prevalence pattern in the Green Spain region. While the prevalence of asthma in adolescents continued to be high in the GAN centres located in this region, such as those in Bilbao, A Coruña and Cantabria (19%, 16.5% and 15.4% compared to 12.8%, 15.2% and 16.7% in the ISAAC-III), other centres, like those in Pamplona and Salamanca, also turned out to have a high prevalence (16.9% and 14.7%, respectively). Cartagena, the only centre in the Mediterranean region, reported a prevalence in adolescents similar to the past one (10.2% compared to 9.9% in the ISAAC-III), which could be indicative of a stabilization in this region in association with differences in climate and the hours of sun exposure.¹⁴

The geographical variation observed in asthma in the ISAAC-III in children aged 6–7 years could not be discerned as clearly in the GAN study. Thus, the prevalence observed in GAN centres located in Green Spain (of 10.9%, 11% and 11.4% in Bilbao, A Coruña and Cantabria, respectively) did not differ from the prevalence found in the only GAN centre in the Mediterranean region (Cartagena, with a prevalence that was only slightly greater at 11.7%). However, these four centres near the coast do differ from centres located inland and at higher altitudes, such as Pamplona and Salamanca, which reported lower figures of 7% and 9.1%, respectively. This coast-versus-inland pattern observed in schoolchildren cannot be generalised to the entire country, nor the epidemiological factors at play in the difference identified, since the study did not include data for

Table 2 Prevalence of asthma symptoms^a. Paper-based GAN questionnaire in schoolchildren aged 6–7 years.

Centre	Sample size	Ever had wheezing	Recent wheezing	Recent wheezing attacks			Recent awakenings		Wheezing limiting speech	Ever had asthma	Recent wheezing during exercise	Recent cough at night
				1–3	4–12	>12	<1 night/wk	>1 night/wk				
Bilbao	2707	1013 (37.4%)	295 (10.9%)	203 (7.5%)	63 (2.3%)	18 (0.7%)	114 (4.2%)	42 (1.6%)	58 (2.1%)	615 (22.7%)	174 (6.4%)	752 (27.8%)
Cantabria	2841	1035 (36.4%)	324 (11.4%)	227 (8.0%)	64 (2.3%)	20 (0.7%)	128 (4.5%)	43 (1.5%)	65 (2.3%)	490 (17.3%)	162 (5.7%)	851 (30.0%)
Cartagena	3509	1304 (37.2%)	411 (117%)	306 (8.7%)	59 (1.7%)	24 (0.7%)	170 (4.8%)	75 (2.1%)	86 (2.5%)	362 (10.3%)	213 (6.1%)	960 (27.4%)
A Coruña	3407	1206 (35.4%)	374 (11.0%)	250 (7.3%)	90 (2.6%)	20 (0.6%)	151 (4.4%)	45 (1.3%)	77 (2.3%)	332 (9.7%)	166 (4.9%)	1054 (30.9%)
Pamplona	2363	680 (28.8%)	165 (6.98%)	151 (6.4%)	25 (1.1%)	1 (0.04%)	87 (3.7%)	20 (0.85%)	26 (1.1%)	150 (6.4%)	78 (3.3%)	525 (22.2%)
Salamanca	2388	797 (33.4%)	219 (9.17%)	154 (6.5%)	41 (1.7%)	7 (0.3%)	78 (3.3%)	35 (1.5%)	27 (1.1%)	191 (8.0%)	103 (4.3%)	544 (22.8%)
Total	17 215	6035 (35.1%)	1788 (10.4%)	1291 (7.5%)	342 (2.0%)	90 (0.5%)	728 (4.3%)	260 (1.5%)	339 (2.0%)	2140 (12.4%)	896 (5.2%)	4686 (27.2%)

^a Prevalence expressed as absolute frequency and percentage, n (%).

Table 3 Prevalence of asthma symptoms^a. Paper-based GAN questionnaire in adolescents aged 13–14 years.

Centre	Sample size	Ever had wheezing	Recent wheezing	Recent wheezing attacks			Recent awakenings		Wheezing limiting speech	Ever had asthma	Recent wheezing during exercise	Recent cough at night
				1–3	4–12	>12	<1 night/wk	>1 night/wk				
Bilbao	3379	1043 (30.9%)	641 (19.0%)	367 (10.9%)	171 (5.1%)	65 (1.9%)	207 (6.1%)	69 (2.0%)	231 (6.8%)	1010 (29.9%)	910 (26.9%)	1194 (35.3%)
Cantabria	4382	1090 (24.9%)	676 (15.4%)	423 (9.7%)	159 (3.6%)	85 (1.9%)	217 (5.0%)	55 (1.3%)	190 (4.3%)	1033 (23.6%)	981 (22.4%)	1378 (31.5%)
Cartagena	3437	657 (19.1%)	350 (10.2%)	217 (6.3%)	65 (1.9%)	25 (0.7%)	108 (3.1%)	35 (1.0%)	105 (3.1%)	513 (14.9%)	477 (13.9%)	810 (23.6%)
A Coruña	3462	813 (23.5%)	570 (16.5%)	324 (9.4%)	144 (4.2%)	65 (1.9%)	170 (4.9%)	50 (1.4%)	137 (4.0%)	712 (20.6%)	737 (21.3%)	1236 (35.7%)
Pamplona	1798	537 (29.9%)	304 (16.9%)	229 (12.7%)	70 (3.9%)	11 (0.6%)	114 (6.3%)	19 (1.1%)	63 (3.5%)	323 (18.0%)	391 (21.8%)	561 (31.2%)
Salamanca	3485	896 (25.7%)	511 (14.7%)	343 (9.8%)	108 (3.1%)	37 (1.1%)	181 (5.2%)	36 (1.0%)	137 (3.9%)	665 (19.1%)	825 (23.7%)	997 (28.6%)
Total	19943	5036 (25.3%)	3052 (15.3%)	1903 (9.5%)	717 (3.6%)	288 (1.4%)	997 (5.0%)	264 (1.3%)	863 (4.3%)	4256 (21.3%)	4321 (21.67%)	6176 (31.0%)

^a Prevalence expressed as absolute frequency and percentage, n (%).

large regions, such as Andalusia, Catalonia, Madrid, the Balearic Islands or the Canary Islands. Other studies conducted in Spain, such as the International Study of Wheezing in Infants, have not found geographical differences in prevalence, with wheezing found in at least one third of infants under study independently of location,^{20,21} although it is well known that wheezing in infants differs from wheezing in school-age children and adolescents.

There is controversy regarding the global trend in asthma prevalence. In the ISAAC-III, some centres with a low asthma prevalence reported an increasing trend, while others with a high baseline prevalence reported decreases, especially in English-speaking countries and Western Europe.¹ In the United States, the prevalence of asthma in the population aged less than 18 years increased from 8.7% to 9.4% in the 2001–2010 period, followed by a plateau until 2013 and a subsequent decrease to 7% by 2019.^{22–24} Data from the National Health Interview Survey of the United States also showed a prevalence peak of 10.9% at age 12–14 years. In Thailand, the data shows that the prevalence of asthma has stabilized in schoolchildren aged 6–7 years (14.6%), with a slight decrease in adolescents (12.5%).²⁵ Other recent data show that the prevalence of asthma is currently stable in Finnish youth.²⁶ The data from our study show that the prevalence of asthma is stabilizing in schoolchildren and increasing in adolescents, although the prevalence remains high in GAN centres that had previously reported a high prevalence in the ISAAC-III. There is likely a potential maximum prevalence in each population that could be reached in specific areas when all predisposed individuals exhibit symptoms, which could explain the asthma prevalence trends in different regions and countries.²⁷

Exercise-induced asthma symptoms were frequent in adolescents, without differences in prevalence between the paper-based and video questionnaires (20.8% vs 21.6%). However, we ought to highlight the low prevalence in children aged 6–7 years based on parental reports (5.2% compared to 5.1% in the ISAAC-III), possibly due to a decreased frequency and intensity of physical activity and greater time spent playing at that age.^{10,11} We did not find a substantial difference in the preva-

lence of nocturnal dry cough in the past year obtained through the paper-based questionnaire (27.2% in schoolchildren versus 31% in adolescents), although it was greater compared to the ISAAC-III (18.9% and 23.1%, respectively).

On the other hand, a past history of asthma was reported in 21.3% of adolescents and 12.4% of schoolchildren, which, compared to the data of the ISAAC-III (14.3% and 11.8%, respectively) corroborates the significant increase in asthma in adolescents (with increases varying from 29.9% in Bilbao to 14.9% in Cartagena) and the stabilization of its prevalence in schoolchildren. The self-reporting of asthma indicates a medical diagnosis, which confirms the need of guaranteeing correct diagnosis and treatment of asthma, especially at a crucial stage like adolescence.

In our study, the comparison by sex revealed a greater prevalence in boys and female adolescents, with differences that were more frequently significant compared to the ISAAC-III. This finding was consistent with those of the GAN centres in Chile and Mexico, where female sex was found to be a significant risk factor for development of asthma in adolescence and female adolescents exhibited the highest prevalence of asthma, while the prevalence in the 6–7 years age group was greater in boys.^{28,29} The observed differences by sex are not clearly understood. Studies in animals show that oestrogens increase Th2-mediated airway inflammation and, on the other hand, in the early years of life, boys exhibit a greater disproportion than girls in the size and growth of the airways compared to the rest of the lung parenchyma.³⁰

When we analysed severe wheezing, defined as the presence in the past year of 4 or more whistling attacks, 1 or more awakenings a week due to wheezing or any wheezing attack that impeded speech, we found a significant increase in prevalence in adolescents (7% in the GAN study vs 5.1% in the ISAAC-III).^{10,11} All of the above, in addition to 5% of adolescents reporting 4 or more asthma exacerbations in the past year, reflect poor asthma control which, combined with the poor adherence characteristic of adolescence, evinces a pressing need for asthma education programmes to achieve adequate self-management of the disease.

Table 4 Prevalence of asthma symptoms^a. Video GAN questionnaire in adolescents aged 13–14 years.

Centre	Sample size	Scene 1			Scene 2			Scene 3			Scene 4			Scene 5		
		Wheezing			Exercise wheezing			Sleep disturbance			Night cough			Severe asthma attack		
		Ever	12 m	1 m	Ever	12 m	1 m	Ever	12 m	1 m	Ever	12 m	1 m	Ever	12 m	1 m
Bilbao	3379	990 (29.3%)	581 (17.2%)	268 (7.9%)	1294 (38.3%)	884 (26.2%)	457 (13.5%)	505 (15.0%)	248 (7.3%)	121 (3.6%)	1287 (38.1%)	851 (25.2%)	350 (10.4%)	658 (19.5%)	408 (12.1%)	189 (5.6%)
Cantabria	4382	1000 (22.8%)	625 (14.3%)	280 (6.4%)	1358 (31.0%)	984 (22.5%)	455 (10.4%)	544 (12.4%)	313 (7.1%)	142 (3.2%)	1279 (29.2%)	870 (19.9%)	305 (7.0%)	531 (12.1%)	345 (7.9%)	163 (3.7%)
Cartagena	3437	478 (13.9%)	256 (7.5%)	113 (3.3%)	616 (17.9%)	390 (11.4%)	181 (5.3%)	257 (7.5%)	131 (3.8%)	50 (1.5%)	981 (28.5%)	580 (16.9%)	189 (5.5%)	314 (9.1%)	182 (5.3%)	78 (2.3%)
A Coruña	3462	778 (22.5%)	498 (14.4%)	193 (5.6%)	1129 (32.6%)	850 (24.6%)	399 (11.5%)	416 (12.0%)	234 (6.8%)	84 (2.4%)	1329 (38.4%)	893 (25.8%)	260 (7.5%)	520 (15.0%)	327 (9.5%)	121 (3.5%)
Pamplona	1798	388 (21.6%)	215 (12.0%)	74 (4.1%)	510 (28.4%)	383 (21.3%)	140 (7.8%)	209 (11.6%)	106 (5.9%)	46 (2.6%)	822 (45.7%)	538 (29.9%)	136 (7.6%)	308 (17.1%)	179 (10.0%)	82 (4.6%)
Salamanca	3485	721 (20.7%)	394 (11.3%)	164 (4.7%)	1000 (28.7%)	657 (18.9%)	300 (8.6%)	345 (9.9%)	155 (4.5%)	66 (1.9%)	1022 (29.3%)	636 (18.3%)	200 (5.7%)	480 (13.8%)	262 (7.5%)	120 (3.4%)
Total	19 943	4355 (21.8%)	2569 (12.9%)	1092 (5.5%)	5907 (29.6%)	4148 (20.8%)	1932 (9.7%)	2276 (11.4%)	1187 (6.0%)	509 (2.6%)	6720 (33.7%)	4368 (21.9%)	1440 (7.2%)	2811 (14.1%)	1703 (8.5%)	753 (3.8%)

^a Prevalence expressed as absolute frequency and percentage, n (%).

Table 5 Comparison of the prevalence^a of asthma symptoms by sex in schoolchildren aged 6 to 7 years and adolescents aged 13 to 14 years in the ISAAC-III versus GAN study (written questionnaire).

Centre	Recent wheezing				Severe wheezing				Ever had asthma			
	ISAAC-III		GAN		ISAAC-III		GAN		ISAAC-III		GAN	
	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male
<i>Bilbao</i>												
6– 7 years	9.9 (8.5–11.5)	14.6 (12.8–16.5)	8.9 (7.4–10.5)	13.0 (11.2–14.9)	3.4 (2.6–4.5)	4.2 (3.3–5.4)	3.6 (2.7–4.8)	4.6 (3.5–5.8)	16.9 (15.1–18.8)	24.6 (22.5–26.9)	19.4 (17.3–21.6)	26.1 (23.8–28.6)
13– 14 years	12.9 (11.3–14.7)	12.6 (10.9–14.5)	20.6 (18.7–22.7)	17.3 (15.5–19.2)	4.9 (3.9–6.1)	5.7 (4.6–7.1)	11.1 (9.6–12.8)	8.1 (6.9–9.6)	19.3 (17.3–21.3)	23.1 (20.9–25.5)	26.2 (24.1–28.5)	33.3 (31.0–35.6)
<i>Cantabria</i>												
6–7 years	NA	NA	10.6 (9.1–12.3)	12.3 (10.7–14.2)	NA	NA	3.5 (2.6–4.6)	5.4 (4.3–6.7)	NA	NA	14.4 (12.6–16.3)	20.1 (18.0–22.3)
13– 14 years	17.6 (15.2–20.2)	15.7 (13.4–18.3)	17.2 (15.7–18.9)	13.3 (11.8–14.8)	8.6 (6.9–10.6)	8.1 (6.4–10.1)	9.0 (7.8–10.2)	5.5 (4.6–6.6)	15.6 (13.3–18.1)	17.9 (15.4–20.5)	23.0 (21.3–24.9)	24.0 (22.2–25.9)
<i>Cartagena</i>												
6– 7 years	9.4 (7.9–11.1)	12.7 (10.9–14.5)	10.2 (8.8–11.7)	13.2 (11.7–14.9)	3.5 (2.6–4.7)	4.6 (3.5–5.8)	4.3 (3.4–5.4)	4.4 (3.5–5.4)	8.7 (7.2–10.3)	12.9 (11.1–14.8)	8.3 (7.1–9.7)	12.3 (10.8–13.9)
13– 14 years	8.6 (7.2–10.2)	11.2 (9.6–13.0)	10.7 (9.3–12.3)	9.3 (7.9–10.8)	2.8 (2.0–3.8)	4.0 (3.0–5.2)	4.8 (3.8–5.9)	3.2 (2.4–4.2)	9.9 (8.4–11.6)	12.8 (11.1–14.6)	14.5 (12.9–16.3)	15.1 (13.4–16.9)
<i>A Coruña</i>												
6– 7 years	12.0 (10.4–13.7)	13.9 (12.1–15.7)	9.7 (8.3–11.2)	12.3 (10.8–14.0)	4.5 (3.5–5.7)	4.8 (3.8–6.0)	4.3 (3.4–5.4)	4.6 (3.7–5.7)	11.6 (10.0–13.3)	16.0 (14.1–17.9)	9.0 (7.6–10.4)	10.6 (9.2–12.2)
13– 14 years	15.0 (13.2–17.0)	15.4 (13.6–17.3)	18.0 (16.2–19.9)	15.1 (13.4–16.9)	6.4 (5.2–7.8)	6.4 (5.2–7.7)	8.7 (7.4–10.2)	7.6 (6.4–8.9)	14.4 (12.6–16.3)	22.3 (20.3–24.5)	19.1 (17.3–21.1)	22.1 (20.1–24.1)
<i>Pamplona</i>												
6– 7 years	6.7 (5.5–8.0)	7.5 (6.2–8.9)	5.9 (4.6–7.4)	8.0 (6.6–9.7)	2.1 (1.4–2.9)	2.5 (1.8–3.4)	1.9 (1.2–2.9)	2.4 (1.6–3.4)	8.1 (6.8–9.5)	11.3 (9.8–12.9)	5.3 (4.1–6.8)	7.3 (5.9–8.9)
13– 14 years	8.2 (6.8–9.8)	7.8 (6.5–9.3)	17.9 (15.5–20.4)	15.8 (13.4–18.4)	3.1 (2.3–4.2)	3.1 (2.3–4.1)	6.9 (5.4–8.7)	6.4 (4.8–8.3)	8.8 (7.4–10.5)	12.7 (11.1–14.4)	17.9 (15.5–20.4)	18.1 (15.5–20.9)
<i>Salamanca</i>												
6– 7 years	NA	NA	6.5 (5.1–8.0)	11.9 (10.1–13.9)	NA	NA	2.4 (1.6–3.4)	3.5 (2.5–4.7)	NA	NA	5.5 (4.2–6.9)	10.5 (8.8–12.4)
13– 14 years	NA	NA	16.6 (14.9–18.4)	12.6 (11.1–14.3)	NA	NA	7.3 (6.1–8.6)	4.6 (3.6–5.7)	NA	NA	18.3 (16.5–20.2)	19.8 (17.9–21.8)

NA, not available.

^a Prevalence expressed as percentage with the corresponding 95% confidence interval in parentheses.

Table 6 Comparison of the prevalence^a of asthma symptoms in schoolchildren aged 6–7 years and adolescents aged 13–14 years in the ISAAC-III versus GAN study (written questionnaire).

Variable	Centre	Children aged 6–7 years				Adolescents aged 13–14 years			
		ISAAC-III		GAN		ISAAC-III		GAN	
		n	Prevalence % (95% CI)	n	Prevalence % (95% CI)	n	Prevalence % (95% CI)	n	Prevalence % (95% CI)
Recent wheezing	Bilbao	371	12.2 (11.0–13.4)	295	10.9 (9.7–12.1)	368	12.8 (11.6–14.0)	641	19 (17.7–20.3)
	Cantabria	NA	NA	324	11.4 (10.3–12.6)	302	16.7 (15.0–18.5)	676	15.4 (14.4–16.5)
	Cartagena	300	11.0 (9.9–12.2)	411	11.7 (10.7–12.8)	283	9.9 (8.8–11.1)	350	10.2 (9.2–11.2)
	A Coruña	389	12.9 (11.7–14.1)	374	11.0 (9.9–12.1)	453	15.2 (13.9–16.6)	570	16.5 (15.2–17.7)
	Pamplona	225	7.1 (6.2–8.0)	165	7.0 (6.0–8.1)	235	8.0 (7.1–9.1)	304	16.9 (15.2–18.7)
	Salamanca	NA	NA	219	9.2 (8.0–10.4)	NA	NA	511	14.7 (13.5–15.9)
	Total	1285	10.7 (10.2–11.3)	1788	10.4 (9.9–10.9)	1641	12.2 (11.6–12.8)	3052	15.3 (14.8–15.8)
Severe wheezing	Bilbao	116	3.8 (3.2–4.6)	111	4.1 (3.4–4.9)	153	5.3 (4.5–6.2)	326	9.6 (8.7–10.7)
	Cantabria	NA	NA	125	4.4 (3.7–5.2)	152	8.4 (7.1–9.8)	323	7.4 (6.6–8.2)
	Cartagena	110	4.0 (3.3–4.8)	153	4.4 (3.7–5.1)	97	3.4 (2.8–4.1)	141	4.1 (3.5–4.8)
	A Coruña	141	4.7 (3.9–5.5)	151	4.4 (3.8–5.2)	190	6.4 (5.5–7.3)	282	8.1 (7.3–9.1)
	Pamplona	73	2.3 (1.8–2.9)	51	2.2 (1.6–2.8)	92	3.1 (2.5–3.8)	120	6.7 (5.6–7.9)
	Salamanca	NA	NA	70	2.4 (2.3–3.7)	NA	NA	207	5.9 (5.2–6.8)
	Total	440	3.7 (3.3–4.0)	661	3.8 (3.6–4.1)	684	5.1 (4.7–5.5)	1399	7.0 (6.7–7.4)
Ever had asthma	Bilbao	628	20.6 (19.2–22.1)	615	22.7 (21.4–24.6)	609	21.1 (19.7–22.7)	1010	29.9 (29.8–33.0)
	Cantabria	NA	NA	490	17.3 (16.1–18.9)	303	16.7 (15.0–18.6)	1033	23.6 (23.0–25.6)
	Cartagena	293	10.7 (9.6–12.0)	362	10.3 (9.5–11.5)	324	11.3 (10.2–12.6)	513	14.9 (14.2–16.7)
	A Coruña	414	13.7 (12.5–15.0)	332	9.7 (8.9–10.9)	551	18.5 (17.1–20.0)	712	20.6 (19.8–22.6)
	Pamplona	307	9.7 (8.7–10.7)	150	6.4 (5.4–7.4)	319	10.9 (9.8–12.1)	323	18.0 (16.5–20.2)
	Salamanca	NA	NA	191	8.0 (7.0–9.2)	NA	NA	665	19.1 (18.4–21.1)
	Total	1642	13.7 (13.1–14.4)	2140	12.4 (12.1–13.1)	2106	15.6 (15.0–16.3)	4256	21.3 (20.8–21.9)

CI, confidence interval; NA, not available.

^a Prevalence expressed as percentage with the corresponding 95% confidence interval in parentheses.

Although the obtained sample was large and representative enough to reduce selection bias, the fact that some of the centres that participated in the ISAAC did not participate in the GAN study could limit the generalization of the findings on the prevalence of asthma and its temporal trends in Spain. The study had other limitations characteristic of its cross-sectional design, like the inability to establish causal relationships or the use of questionnaires that may give rise to recall bias in parental responses, although limiting the questions to the past 12 months helped reduce this bias. As regards the diagnosis of asthma using data on symptoms obtained through questionnaires as opposed to a medical diagnosis, this is the best feasible approach to compare cities and countries in large-scale epidemiological studies.

In short, the GAN study in Spain, conducted in a large sample of adolescents aged 13–14 years and schoolchildren aged 6–7 years, found a high prevalence of asthma, with an increasing trend in adolescents and stabilization in schoolchildren. The geographical variations detected in 2002 were not as clearly discerned in this study, although it was confirmed that regions where the prevalence of asthma was high in the past continued to have a high prevalence. On the other hand, the observed increases in the prevalence of sleep disturbances, severe asthma and self-reported history of asthma reflect poor asthma control and suggest that asthma has been underdiagnosed and undertreated in recent years.

Funding

This study was funded by research grants from the Instituto de Investigación Sanitaria Valdecilla (IDIVAL) of Cantabria, PRIMVAL 17/01 and 18/01 (Centro Cantabria); the Ministry of Economy and Competitiveness of Spain, the Instituto de Salud Carlos III: Proyectos de Investigación en Salud PI17/00179 (Centro Cartagena), PI17/00756 (Centro Bilbao), PI17/00694 (Centro Pamplona); the Fundación María José Jove (Centro A Coruña); Gerencia Regional de Salud de la Junta de Castilla y León (GRS 1239/b/16) and the Sociedad Española de Inmunología Clínica, Alergología y Asma Pediátrica (Centro Salamanca).

Conflicts of interest

The authors have no conflicts of interest to declare.

Acknowledgments

We thank the collaboration of the Departments of Education and Departments of Health of the autonomous communities where participating centres are located in Spain, and the cooperation of the teaching staff of primary and secondary schools and parents and students, without whose consent and altruistic help the study would not have been possible.

Appendix A. Members of GAN centres in Spain

Cartagena GAN centre (national coordinating centre): L. García-Marcos, M. Sánchez-Solís, Paediatric Pulmonology and Allergy Unit, Hospital Infantil Universitario Virgen de la Arrixaca, Universidad de Murcia. Instituto Murciano de Investigación Biosanitaria IMIB, Murcia, Spain. A. Martínez-Torres, Paediatric Pulmonology and Allergy Unit and Nursing Research Group, Hospital Infantil Universitario Virgen de la Arrixaca. Instituto Murciano de Investigación Biosanitaria IMIB, Murcia, Spain. V. Pérez-Fernández, E. Morales Bartolomé, Universidad de Murcia. Instituto Murciano de Investigación Biosanitaria IMIB, Murcia, Spain. J.J. Guillén-Pérez, J.F. Amoraga Bernal, J. Llamas Fer-

nández and A. García Coy. Cartagena Public Health System. Department of Health of Murcia, Spain. **Bilbao GAN centre:** C. González Díaz, A. González Hermosa, J. Rementeria Radigales. Department of Paediatrics. Hospital Universitario Basurto, Bilbao, Vizcaya, Spain. **Cantabria GAN centre:** A. Bercedo Sanz, L. Lastra Martínez, R. Pardo Crespo. S. Peñil Sánchez. Servicio Cántabro de Salud. Instituto de Investigación Sanitaria Valdecilla. IDIVAL. Cantabria, Spain. **A Coruña GAN centre:** A. López-Silvarrey Varela, Fundación María José Jove; Servicio Galego de Saúde (SERGAS), A Coruña, Spain. T.R. Pérez Castro, Cardiovascular Research Group (GRINCAR); Cardiovascular Epidemiology, Primary Care and Nursing (INIBIC); School of Nursing and Podiatry, Universidad de A Coruña, Spain. A. Otero Rodríguez, A. Garea Otero, A. Torrado Nogueira, J. Iglesias López, F.J. González Barcala. Servicio Galego de Saúde (SERGAS), A Coruña, Spain. R. Montero López, St. Josef Braunau Hospital, Braunau, Austria. **Pamplona GAN centre:** I. Aguinaga-Ontoso, F. Guillén-Grima, Department of Health Sciences, Universidad Pública de Navarra (UPNA), Pamplona, Navarra, Spain and Instituto de Investigación Sanitaria de Navarra (IdiSNA), Spain. E. Rayón-Valpuesta, J. Coque-Rubio, O. Alvarez-Flames, S. Sola-Cía, R. Saenz-Mendia, R. García-Orellan, X. Elizalde, Department of Health Sciences, Universidad Pública de Navarra (UPNA), Pamplona, Spain. IdiSNA, Instituto de Investigación Sanitaria de Navarra. S. Monje-Ortega, Hospital Santa Marina-Osakidetza, Bilbao, Vizcaya. **Salamanca GAN centre:** J. Pellegrini Belinchón, Centro de Salud Pizarrales, Salamanca, Spain. Department of Biomedical Sciences and Diagnosis, Universidad de Salamanca, Spain. S. Arriba-Méndez, Department of Paediatrics, Hospital Universitario Salamanca and Instituto de Investigación Biomédica de Salamanca IBSAL, Spain. A. Marín-Cassinello, Department of Paediatric Pulmonology and Allergy, Hospital General Universitario Santa Lucía, Cartagena, Murcia, Spain. M. Domínguez, Centro de Salud San Juan, Salamanca, Spain. MC. Sánchez-Jiménez, Centro de Salud Tejares, Salamanca and Universidad de Salamanca, Spain. M. M. López-González, Centro de Salud Pizarrales, Salamanca, Spain. M.C. Vega-Hernández, Department of Statistics, Universidad de Salamanca, Spain. M. Polo-De Dios. Centro de Salud Zamora Sur. Zamora, Spain.

References

1. Asher MI, Montefort S, Björkstén B, Lai CKW, Strachan DP, Weiland SK, et al. and the ISAAC Phase Three Study group. Worldwide variation in prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and atopic eczema: ISAAC. The International Study of Asthma and Allergies in Childhood (ISAAC) Steering Committee. *Lancet*. 1998;351:1225–32.
2. Mallol J, Crane J, von Mutius E, Odhiambo J, Keil U, Stewart A, et al. The International Study of Asthma and Allergies in Childhood (ISAAC) Phase Three: a global synthesis. *Allergol Immunopathol (Madr)*. 2013;41:73–85.
3. Global Burden of Disease Study 2019 (GBD 2019) data resources [accessed 18 Jul 2021]. Available from: <http://www.healthdata.org>.
4. Genuneit J, Cantelmo JL, Weinmayr G, Wong GWK, Cooper PJ, Riiikjær MA, et al. A multi-centre study of candidate genes for wheeze and allergy: the International Study of Asthma and Allergies in Childhood Phase 2. *Clin Exp Allergy*. 2009;39:1875–88.
5. Weinmayr G, Keller F, Kleiner A, du Prel JB, Garcia-Marcos L, Batllés-Garrido J, et al. Asthma phenotypes identified by latent class analysis in the ISAAC phase II Spain study. *Clin Exp Allergy*. 2013;43:223–32.
6. García-Marcos L, Miner Canflanca I, Batllés Garrido JB, López-Silvarrey Varela A, García-Hernández G, Guillén Grima F, et al. Relationship of asthma and rhinoconjunctivitis with obesity,

- exercise and Mediterranean diet in Spanish schoolchildren. *Thorax*. 2007;62:503–8.
7. Nagel G, Weinmayr G, Kleiner A, García-Marcos L, Strachan DP. Effect of diet on asthma and allergic sensitisation in the International Study on Allergies and Asthma in Childhood (ISAAC) Phase Two. *Thorax*. 2010;65:516–22.
 8. García-Marcos L, Castro-Rodriguez JA, Weinmayr G, Panagiotakos DB, Priftis KN, Nagel G. Influence of Mediterranean diet on asthma in children: a systematic review and meta-analysis. *Pediatr Allergy Immunol*. 2013;24:330–8.
 9. Mitchell EA, Beasley R, Bjorksten B, Crane J, García-Marcos L, Keil U, et al. The association between BMI, vigorous physical activity and television viewing and the risk of symptoms of asthma, rhinoconjunctivitis and eczema in children and adolescents: ISAAC Phase Three. *Clin Exp Allergy*. 2013;43:73–84.
 10. García-Marcos L, Blanco Quirós A, García Hernández G, Guillén-Grima F, González Díaz C, Carvajal Uruña I, et al. Stabilization of asthma prevalence among adolescents and increase among schoolchildren (ISAAC phases I and III) in Spain. *Allergy*. 2004;59:1301–7.
 11. Carvajal-Uruña I, García-Marcos L, Busquets-Monge R, Morales Suarez-Varela M, García de Andoin N, Batlles-Garrido J, et al. Geographic variation in the prevalence of symptoms in Spanish children and adolescents. *International Study of Asthma and Allergies in Childhood (ISAAC) Phase 3, Spain*. *Arch Bronconeumol*. 2005;41:659–66.
 12. Bercedo Sanz A, Redondo Figuero C, Lastra Martínez L, Gómez Serrano M, Mora González E, Pacheco Cumani M, et al. Prevalencia de asma bronquial, rinitis alérgica y dermatitis atópica en adolescentes de 13-14 años en Cantabria. *Bol Pediatr*. 2004;44:9–19.
 13. García-Marcos L, Batlles-Garrido J, Blanco-Quiros A, García Hernández G, Guillén-Grima F, González Díaz C, et al. Influence of two different geo-climatic zones on the prevalence and time trends of asthma symptoms among Spanish adolescents and schoolchildren. *Int J Biometeorol*. 2009;53:53–60.
 14. Arnedo-Pena A, García-Marcos L, Bercedo-Sanz A, Aguinaga-Ontoso I, González-Díaz C, García-Merino A, et al. Prevalence of asthma symptoms in schoolchildren, and climate in west European countries: an ecologic study. *Int J Biometeorol*. 2013;57:775–84.
 15. García-Marcos L, Morales Suarez-Varela M, Miner Canflanca I, Batllés Garrido J, Blanco Quirós A, López Sivarrey Varela A, et al. BCG immunization at birth and atopic diseases in a homogeneous population of Spanish schoolchildren. *Int Arch Allergy Immunol*. 2005;137:303–9.
 16. Arnedo-Pena A, García-Marcos L, Carvajal Uruña I, Busquets Monge R, Morales Suarez-Varela M, Miner Canflanca I, et al. [Air pollution and recent symptoms of asthma, allergic rhinitis, and atopic eczema in schoolchildren aged between 6 and 7 years]. *Arch Bronconeumol*. 2009;45:224–9.
 17. Ellwood P, Asher MI, Billo NE, Bissell K, Chiang CY, Ellwood EM, et al. The Global Asthma Network rationale and methods for Phase I global surveillance: prevalence, severity, management and risk factors. *Eur Respir J*. 2017;49:1601–5.
 18. Ellwood P, Ellwood E, Rutter C, Perez-Fernandez V, Morales E, García-Marcos L, et al. and on behalf of the GAN Phase I Study Group. Global Asthma Network Phase I Surveillance: geographical coverage and response rates. *J Clin Med*. 2020;9:3688, <http://dx.doi.org/10.3390/jcm9113688>.
 19. Ellwood P, Williams H, Ait-Khaled N, Bjorksten B, Robertson C, Group IPIS. Translation of questions: the International Study of Asthma and Allergies in Childhood (ISAAC) experience. *Int J Tuberc Lung Dis*. 2009;13:1174–82.
 20. Bercedo-Sanz A, Lastra-Martinez L, Pellegrini-Belinchon J, Vicente-Galindo E, Lorente-Toledano F, Garcia-Marcos L. Wheezing and risk factors in the first year of life in Cantabria, Spain. *The EISL study*. *Allergol Immunopathol (Madr)*. 2015;43:543–52.
 21. Pellegrini-Belinchon J, Miguel-Miguel G, Dios-Martin B, Vicente-Galindo E, Lorente-Toledano F, Garcia-Marcos L. Study of wheezing and its risk factors in the first year of life in the Province of Salamanca, Spain. *The EISL Study*. *Allergol Immunopathol (Madr)*. 2012;40:164–71.
 22. Akinbami LJ, Simon AE, Rossen LM. Changing trends in asthma prevalence among children. *Pediatrics*. 2016;137:1–7.
 23. Zahran HS, Bailey CM, Damon SA, Garbe PL, Breyse PN. Vital signs: asthma in children-United States, 2001-2016. *MMWR Morb Mortal Wkly Rep*. 2018;67:149–55.
 24. Centers for Disease Control and Prevention. Most recent national asthma data [accessed 18 Jul 2021]. Available from: https://www.cdc.gov/asthma/most_recent_national_asthma_data.htm.
 25. Chinratapisit S, Suratannon N, Pacharn P, Sritipsukho P, Vichyanond P. Prevalence and severity of asthma, rhinoconjunctivitis and eczema in children from the Bangkok area: The Global Asthma Network (GAN) Phase I. *Asian Pac J Allergy Immunol*. 2019;37:226–31.
 26. Reijula J, Latvala J, Mäkelä M, Siitonen S, Saario M, Hahtela T. Long-term trends of asthma, allergic rhinitis and atopic eczema in Young Finnish men: a retrospective analysis, 1926-2017. *Eur Respir J*. 2020;56:1902144, <http://dx.doi.org/10.1183/13993003.02114-2019>.
 27. García-Marcos L. Asthma prevalence trends in Chile: a prelude of the Global asthma Network findings? *Allergol Immunopathol (Madr)*. 2019;47:311–2.
 28. Mallol J, Aguirre V, Mallol-Simmonds M, Matamala-Bezmalinovic A, Calderón-Rodríguez L, Osses-Vergara F. Changes in the prevalence of asthma and related risk factors in adolescents: three surveys between 1994 and 2015. *Allergol Immunopathol (Madr)*. 2019;47:313–21.
 29. Del-Río-Navarro BE, Berber A, Reyes-Noriega N, Navarrete-Rodríguez EM, García-Almaráz R, Mérida-Palacio JV, et al. Have asthma symptoms in Mexico changed in the past 15 years? Time trends from the International Study of Asthma and Allergies in Childhood to the Global Asthma Network. *Allergol Immunopathol (Madr)*. 2021;49:1–10.
 30. Fuseini H, Newcomb DC. Mechanisms driving gender differences in asthma. *Curr Allergy Asthma Rep*. 2017;17:19, <http://dx.doi.org/10.1007/s11882-017-0686-1>.