

SCIENTIFIC LETTERS

Growth and pubertal development in girls adopted from China: 15-year follow-up



Crecimiento y desarrollo puberal en niñas adoptadas de China: seguimiento de 15 años

Dear Editor,

Prospective studies on the physical development of girls adopted from China to date, to our knowledge, have had follow-up periods of less than 4 years.^{1–3} The purpose of this scientific letter is to contribute data on the temporal trends in weight, height, body mass index (BMI) and head circumference in 30 Chinese girls adopted by Spanish families between 2001 and 2007 who were followed up for 15 years from the post-arrival medical evaluation, the age at thelarche (Tanner stage II of pubertal development) and the age at menarche.

Follow-up evaluations and measurements took place every 6 months and were always performed by the same clinician. Families were instructed on how to watch for thelarche and menarche and to schedule an appointment once they happened (in these particular visits, the percentage body fat was calculated with the Deurenberg

equation). We entered data for the study variables in an Excel spreadsheet, also used to calculate the mean and standard deviation (SD). Using the WHO Anthroll (0–5 years) and WHO AnthroPlus (5–19 years) applications, we compared the anthropometric measurements obtained in the sample with the World Health Organization (WHO) growth standards, calculating the corresponding z scores and percentiles (PCTL) for age and sex (note: these applications do not calculate the z score or percentile of the head circumference from age 5 years or of the weight from age 10 years). We defined significant catch-up growth for any given variable as a z score increase of +0.5 SDs or greater.

Table 1 presents the anthropometric measurements obtained in the initial evaluation and in the annual visits over the 15-year follow-up, comparing them with the WHO growth standards. Fig. 1 shows the temporal trends in the height and BMI z scores using the WHO growth standards as reference.

The mean weight and height z scores in the post-arrival evaluation were very similar to those reported by Miller and Hendrie⁴ in girls adopted from China at a mean age of 15.7 months. We observed significant catch-up growth in weight and BMI in the first year following the post-arrival evaluation, in height in the second year, and in head circumference in the third year. In a study conducted 3 years after the




Figure 1 Height and body mass index (BMI) z scores in girls adopted from China (n = 30) over a 15-year period. Comparison with the World Health Organization child growth standards.

Table 1 Weight, height, body mass index and head circumference in girls adopted from China (n = 30) in the initial post-arrival evaluation and during the 15-year follow-up. Comparison with the World Health Organization child growth standards.

	Weight (kg)				Height (cm)				BMI			Head circumference (cm)				
	Mean	±SD	z	PCTL	Mean	±SD	z	PCTL	Mean	z	PCTL	Mean	±SD	z	PCTL	
<i>Post-arrival evaluation</i>																
Age, mean ± SD: 16 ± 5 months	8.53	±1.65	-1.16	12.3	74.45	±6.91	-1.49	6.8	15.4	-0.38	35.2	44.37	±1.52	-1.09	13.9	
<i>Age/follow-up</i>																
2 years 4 months/+1 year	11.81	±1.46	-0.46	32.3	86.94	±5.64	-1.05	14.8	15.9	0.24	59.7	46.74	±0.99	-0.77	22.1	
3 years 4 months/+2 years	13.72	±1.65	-0.57	28.5	95.18	±5.94	-0.97	16.6	15.4	0.03	51.4	47.98	±0.88	-0.64	26.0	
4 years 4 months/+3 years	15.83	±2.22	-0.49	31.1	102.41	±6.03	-0.87	19.3	15.3	0.04	51.4	48.79	±0.90	-0.57	28.5	
5 years 4 months/+4 years	17.73	±2.87	-0.47	32.0	108.65	±6.66	-0.61	26.9	15.0	-0.15	44.0	49.23	±1.02			
6 years 4 months/+5 years	20.20	±3.60	-0.27	39.4	115.47	±6.72	-0.38	35.3	15.2	-0.10	46.0	49.71	±1.04			
7 years 4 months/+6 years	22.59	±4.06	-0.23	41.1	121.84	±6.94	-0.23	40.7	15.2	-0.17	43.3	50.38	±1.17			
8 years 4 months/+7 years	24.92	±4.90	-0.33	37.2	127.34	±7.33	-0.27	39.2	15.4	-0.27	39.5	50.85	±1.24			
9 years 4 months/+8 years	26.62	±4.66	-0.65	25.7	130.76	±5.70	-0.68	24.9	15.6	-0.39	34.7	51.34	±1.30			
10 years 4 months/+9 years	29.38	±3.75			136.63	±4.87	-0.70	24.0	15.7	-0.58	28.1	51.95	±1.45			
11 years 4 months/+10 years	34.11	±4.28			143.09	±5.58	-0.67	25.1	16.7	-0.42	33.9	52.51	±1.51			
12 years 4 months/+11 years	38.43	±5.41			148.12	±4.94	-0.79	21.6	17.5	-0.35	36.3	53.17	±1.79			
13 years 4 months/+12 years	43.19	±5.54			152.64	±4.62	-0.77	22.1	18.5	-0.23	40.7	53.86	±1.69			
14 years 4 months/+13 years	48.50	±3.81			155.17	±5.22	-0.80	21.1	20.1	0.11	54.2	54.64	±1.67			
15 years 4 months/+14 years	51.33	±5.35			157.04	±5.26	-0.74	22.9	21.1	0.23	59.1	55.13	±1.75			
16 years 4 months/+15 years	52.06	±4.73			157.30	±5.08	-0.80	21.2	20.7	-0.04	48.5	55.15	±1.74			

BMI, body mass index; PCTL, percentile; SD, standard deviation; z, z score.

post-arrival evaluation in girls adopted from China, Palacios et al.³ found significant catch-up growth in every anthropometric parameter. In the longitudinal analysis of the height z score, we found catch-up growth through age 8 years, with a decrease between ages 8 and 10 years and stabilization from 10 to 16 years. In the longitudinal analysis of the BMI z score, compared to linear growth, we found a greater catch-up growth through age 6 years, similar growth between ages 6 and 10 years, and progressive growth between ages 10 and 16 years.

The mean age at thelarche was 10 years and 1 month (SD, 1 year and 3 months). At this follow-up visit, the mean weight was 30.3 kg (SD, 4.6), the mean height 137.1 cm (SD, 5.8; z, -0.40; 32.9th PCTL), the mean BMI 16.1 (SD, 1.7; z, -0.30; 38.3th PCTL) and the mean percentage body fat 16.1% (SD, 1.9%). The mean age at menarche was 12 years and 2 months (SD, 1 year). At this visit, the mean weight was 38.7 kg (SD, 3.7), the mean height 149.5 cm (SD, 5.8 cm; z, -0.44; 32.9th PCTL), the mean BMI 17.3 (SD, 1.2; z, -0.38; 35.1th PCTL) and the percentage body fat 18.1% (SD, 1.3%). Hayes and Tan,⁵ in a survey of the adoptive parents of a cohort of 814 girls adopted from China into North America, found a mean age at menarche of 12.37 years (95% CI, 11.84–13 years). In the group of adopted girls analysed in our study, we found a mean linear growth of 20.2 cm from the beginning of Tanner stage II to the time the adult height was achieved; the peak height velocity between thelarche and menarche was 12.4 cm, and the mean remaining linear growth after menarche was 7.8 cm. These data suggest that pubertal development in these girls could be considered normal.⁶ However, the height achieved by age 16 years and 4 months, compared to Chinese linear growth charts and the WHO child growth standards, was smaller by 2.85 cm and 5.4 cm, respectively.⁷

References

1. Oliván Gonzalvo G. Estudio auxológico de niñas adoptadas en China [Auxologic study of girls adopted in China]. *An Pediatr (Barc)*. 2006;65:88–90, <http://dx.doi.org/10.1157/13090906>.
2. Cohen NJ, Lojkasek M, Zadeh ZY, Pugliese M, Kiefer H. Children adopted from China: a prospective study of their growth and development. *J Child Psychol Psychiatry*. 2008;49:458–68, <http://dx.doi.org/10.1111/j.1469-7610.2007.01853.x>.
3. Palacios J, Román M, Camacho C. Growth and development in internationally adopted children: extent and timing of recovery after early adversity. *Child Care Health Dev*. 2011;37:282–8, <http://dx.doi.org/10.1111/j.1365-2214.2010.01142.x>.
4. Miller LC, Hendrie NW. Health of children adopted from China. *Pediatrics*. 2000;105:E76, <http://dx.doi.org/10.1542/peds.105.6.e76>.
5. Hayes P, Tan TX. Timing of menarche in girls adopted from China: a cohort study. *Child Care Health Dev*. 2016;42:859–62, <http://dx.doi.org/10.1111/cch.12393>.
6. Soriano Guillén L. Pubertad normal y variantes de la normalidad. *Pediatr Integral*. 2015;19:380–8.
7. Zong XN, Li H. Construction of a new growth references for China based on urban Chinese children: comparison with the WHO growth standards. *PLoS One*. 2013;8:e59569, <http://dx.doi.org/10.1371/journal.pone.0059569>.

Gonzalo Oliván-Gonzalvo

Centro de Pediatría y Adopción Internacional, Zaragoza, Spain

E-mail address: g.olivan@comz.org

<https://doi.org/10.1016/j.anpede.2023.03.014>
2341-2879/

© 2023 Asociación Española de Pediatría. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Modified Valsalva manoeuvre in paroxysmal supraventricular tachycardia: Case series



Maniobra de Valsalva modificada en taquicardia paroxística supraventricular: serie de casos

Dear Editor,

Paroxysmal supraventricular tachycardia (PSVT) is the most frequent form of sustained tachycardia in childhood. It is defined as tachycardia caused by a rhythm disorder originating above the His bundle.

Its most frequent electrocardiographic presentation is regular tachycardia with a narrow QRS complex. The heart rate is usually greater than 180–220 beats per minute (bpm)

in infants, and at least 120–150 bpm in older children. It responds to vagal manoeuvres and drugs that slow conduction through the atrioventricular node. Clinically, it is well tolerated, but when it is prolonged, patients may develop signs of congestive heart failure.¹

The main goal of acute treatment is to stop the arrhythmia, slow down the ventricular response and restoring the sinus rhythm. In stable patients, the initial management should consist of vagal manoeuvres, as they can succeed in ending the tachycardia in cases involving the atrioventricular node in the conduction pathway. In school-aged children and adolescents, this is mainly achieved through the Valsalva manoeuvre, as both ocular pressure and carotid sinus massage are not recommended in these age groups.² In paediatric patients, the effectiveness of vagal manoeuvres ranges between 30% and 60%.^{3,4}

One of the ways to increase the vagal parasympathetic tone is to increase venous return. Studies in adults have investigated different modifications of the traditional Valsalva manoeuvre technique to increase vagal tone. In particular, the study conducted by Appelboom et al.⁵ found a statistically significant increase in the effectiveness of

DOI of refers to article: <https://doi.org/10.1016/j.anpede.2023.04.007>