



## ORIGINAL ARTICLE

## Socioeconomic and gender inequalities in childhood obesity in Spain



Enrique Gutiérrez-González<sup>a,1</sup>, Francisco Sánchez Arenas<sup>b,1</sup>,  
Ana María López-Sobaler<sup>c,d</sup>, Blanca Andreu Ivorra<sup>e</sup>, Almudena Rollán Gordo<sup>a</sup>,  
Marta García-Solano<sup>a,\*</sup>

<sup>a</sup> Subdirección General de Nutrición, Agencia Española de Seguridad Alimentaria y Nutrición, Ministerio de Consumo, Madrid, Spain

<sup>b</sup> Servicio de Medicina Preventiva, Hospital Universitario de La Princesa, Madrid, Spain

<sup>c</sup> Departamento de Nutrición y Ciencias de los Alimentos, Facultad de Farmacia, Universidad Complutense de Madrid, Madrid, Spain

<sup>d</sup> Grupo de Investigación "Valoración Nutricional de Individuos y Colectivos: Metodología y Aplicaciones" (VALORNUT), Universidad Complutense de Madrid, Madrid, Spain

<sup>e</sup> Servicio de Medicina Preventiva, Hospital Universitario de Alcorcón, Madrid, Spain

Received 16 February 2023; accepted 22 May 2023

Available online 28 July 2023

### KEYWORDS

Obesity;  
Overweight;  
Child;  
Child poverty;  
Socioeconomic  
disparities in health;  
Social determinants  
of health;  
Gender equity;  
Spain

### Abstract

**Introduction:** Childhood obesity is inversely associated with household socioeconomic status in high-income countries. Our aim was to explore the association between childhood weight status and household socioeconomic status in Spain in relation to family characteristics, perceptions and lifestyle habits and the school environment.

**Methods:** We performed a descriptive analysis of child weight status according to family socioeconomic factors and mediating variables based on data from the ALADINO 2019 study in a sample of 16,665 schoolchildren representative of the population aged 6–9 years in Spain.

**Results:** The prevalence of childhood obesity in households with low socioeconomic status (26.8% boys; 20.4% girls) was, in both sexes, twice as high as in those with higher socioeconomic status (12.1% boys; 8.7% girls). Unhealthy eating habits, sedentary lifestyles (mainly in girls) and the presence of screens in the bedroom (more prevalent in boys) were more frequent in school-aged children from low-income households. On the other hand, in the most disadvantaged households, a history of breastfeeding and physical activity (especially in girls) were less frequent. Similarly, schools attended by children from low-income households were less likely to have their own kitchens and indoor gymnasiums or offer sports activities.

**Conclusion:** A lower household socioeconomic status was associated with poorer dietary and physical activity habits and certain characteristics of the family and school environments that mediate the inverse association between household socioeconomic status and the prevalence

DOI of original article: <https://doi.org/10.1016/j.anpedi.2023.05.013>

\* Corresponding author.

E-mail address: [mgarsol@gmail.com](mailto:mgarsol@gmail.com) (M. García-Solano).

<sup>1</sup> Both authors have contributed equally as first authors of the article.

of childhood obesity. Girls were less physically active and reported more sedentary lifestyles, while boys had greater access to screens. Interventions to combat childhood obesity should address the identified inequalities.

© 2023 Asociación Española de Pediatría. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## PALABRAS CLAVE

Obesidad;  
Sobrepeso;  
Infantil;  
Pobreza infantil;  
Desigualdades  
socioeconómicas en  
salud;  
Determinantes  
sociales de la salud;  
Equidad de género;  
España

## Desigualdades socioeconómicas y de género en la obesidad infantil en España

### Resumen

**Introducción:** La obesidad infantil se asocia inversamente con el nivel socioeconómico familiar en países desarrollados. El objetivo es evaluar la asociación entre la situación ponderal infantil y el nivel socioeconómico familiar, según características familiares, percepciones, hábitos de vida y entorno escolar.

**Metodología:** Análisis descriptivo de la situación ponderal infantil según factores socioeconómicos familiares y variables mediadoras en el estudio ALADINO 2019, en una muestra de 16.665 escolares representativa de la población escolar de 6 a 9 años en España.

**Resultados:** La prevalencia de obesidad infantil en hogares de bajo nivel socioeconómico (26,8% niños; 20,4% niñas) era, en ambos sexos, el doble de la de los de mayor nivel (12,1% niños; 8,7% niñas). En escolares de familias de baja renta eran más frecuentes hábitos alimentarios poco saludables, el sedentarismo (principalmente en niñas) y la presencia de pantallas en la habitación (más en niños). Por el contrario, en hogares más desfavorecidos eran menos frecuentes el antecedente de lactancia materna y la práctica de actividad física (especialmente en niñas). La disponibilidad de cocina propia, actividades deportivas y gimnasio cubierto en los centros era también menos habitual para los escolares de familias de menor renta.

**Conclusiones:** Un menor nivel socioeconómico del hogar se asocia con peores hábitos alimentarios y de actividad física y ciertas características del entorno familiar y el escolar, que a su vez median la asociación inversa que existe entre el nivel socioeconómico y la prevalencia de obesidad infantil. Las niñas realizan menos actividad física y presentan más sedentarismo mientras que los niños tienen más acceso a pantallas. Las intervenciones para combatir la obesidad infantil deben contemplar las desigualdades identificadas.

© 2023 Asociación Española de Pediatría. Publicado por Elsevier España, S.L.U. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Childhood overweight and obesity, jointly referred to as excess weight, are associated with multiple disorders and diseases in childhood, such as type 2 diabetes, high blood pressure or metabolic syndrome,<sup>1,2</sup> as well as an increased probability of obesity and associated comorbidities (such as cardiovascular disease and certain forms of cancer) in adulthood.<sup>3</sup>

In recent decades, the prevalence of childhood excess weight has increased in Europe, including Spain.<sup>4</sup> According to the 2018–2020 report of the Childhood Obesity Surveillance Initiative (COSI) of the World Health Organization (WHO) Regional Office for Europe, the prevalence of excess weight in school-aged children (ages 6–9 years) was 31% in boys and 28% in girls, while the prevalence for obesity was 14% in boys and 10% in girls.<sup>5</sup> In Spain, the most recent round of the nationwide study of nutrition, physical activity, child development and obesity (known as ALADINO) of the Agencia Española de Seguridad Alimentaria y Nutrición (Spanish Food Safety and Nutrition Agency, AESAN), conducted in 2019, found a prevalence of excess weight of

41% in boys and 40% in girls and of obesity of 19% in boys and 15% in girls,<sup>6</sup> with a decline and subsequent stabilization of childhood excess weight trends between 2011 and 2019, mainly on account of a decrease in the prevalence of overweight in boys. Given the magnitude and potential repercussions of this problem, it is a major public health challenge.<sup>7</sup>

Childhood obesity results from a complex interaction of factors, many of which are associated with the socioeconomic and cultural context, family background and school environment of the children.<sup>1,7</sup> On the other, child poverty is a multifactorial phenomenon with a negative impact on dimensions as important as physical health, academic performance or emotional wellbeing.<sup>8</sup> The child poverty rate in Spain is among the highest in Europe, with 27% of children and adolescents living in poverty in 2020.<sup>9</sup>

Childhood obesity and socioeconomic status are inversely correlated in high-income countries such as Spain, where childhood obesity is more frequent in families with lower versus higher socioeconomic status,<sup>10</sup> as evinced by the different rounds of the ALADINO study<sup>6,11,12</sup> or the National Health Survey.<sup>13</sup> Yet, the evidence of the factors that medi-

ate the association between childhood obesity and poverty in Spain is scarce.

The aim of our study was to assess the association between weight status in children aged 6–9 years in Spain and household socioeconomic variables from a sex-gender perspective, exploring factors related to household characteristics, lifestyle habits and the school environment that may mediate this association.

## Methods

### Overall design and sample of the ALADINO 2019 study

We obtained the data to conduct the analysis from those collected in the ALADINO 2019 study, a cross-sectional, observational and descriptive study representative of the population of schoolchildren aged 6–9 years attending primary education (PE) schools in Spain. It corresponds to the fifth round of the COSI (the fourth in Spain), the methodology and design of which have been described in previous publications.<sup>11,12</sup>

We selected participants by random multi-stage cluster sampling categorised into 4 age groups (6, 7, 8 and 9 years), enrolled in years 1–4 of primary education. We selected PE schools at random, stratifying by autonomous community (AC) and municipality size, and then selected one classroom per school year per selected school by simple random sampling, including every student in the selected classrooms who attended school on the day of the visit, whose parents or legal guardians signed the informed consent form, and who agreed to undergo the necessary measurements. The resulting sample consisted of 16 665 students, with a uniform sex distribution, from 276 schools. The study was approved by the Ethics Committee of the Hospital Clínico San Carlos of Madrid.

Based on the models and methodological specifications of the COSI-EURO WHO surveys,<sup>14,15</sup> we developed 3 questionnaires to be completed by the parents or legal guardians, the schools and the examiners. The weight, height and waist circumference were measured according to standardised procedures with properly calibrated equipment by qualified examiners.<sup>15,16</sup>

### Study variables

To assess the weight status of the children, we used the body mass index ( $BMI = \text{weight [kg]} / \text{height}^2 [\text{m}^2]$ ) using the WHO growth standards as reference,<sup>17</sup> and the waist-to-height ratio (WtHR), defining central obesity as a WtHR greater than 0.5.<sup>18</sup>

We considered the following socioeconomic variables: household income, parental educational attainment and parental employment. In relation to the family characteristics and history, we collected data on nationality, parental weight status and history of breastfeeding. As for perceptions and lifestyle habits, we analysed the parental perception of the child's weight status, the results of the COSI questionnaire on unhealthy dietary habits,<sup>19</sup> physical activity and sedentary behaviour and the number of screens in the child's bedroom. The variables related to the school

environment were the type of school (public, private or private but publicly funded), availability and type of food services, organization of extracurricular sports activities and availability of indoor sports facilities.

## Analysis of results and statistical methods

Since some ACs increased their respective samples to obtain results representative for their region, there was an increase relative to the allocated proportion in the nationwide sample, so in our analysis we adjusted the results for sex, age and AC.

We conducted a descriptive analysis of the characteristics of the sample, the weight status of the schoolchildren in relation to variables related to the socioeconomic status of the family and of the mediator variables in relation to the income level. We categorised every variable and calculated absolute frequencies and percentages with 95% confidence intervals (CIs). We used the  $\chi^2$  test to analyse the association between qualitative variables, considering differences significant if the *P* value was less than 0.05. We also made a quantitative analysis of the BMI and WtHR variables, calculating the median since the data did not follow a normal distribution, as determined with the Kolmogorov-Smirnov test. The statistical analysis was performed with the Stata package, version 16 (StataCorp LLC, Spain).

## Results

**Table 1** summarises the general characteristics of the sample by sex. Compared to girls, boys had obesity more frequently, were more physically active and had more screens in their room, and their parents were more likely to underestimate their weight status; we did not find differences in any other variables based on sex.

### Association between child weight status and socioeconomic status

The prevalence of obesity and overweight decreased as household income increased, a gradient that was more pronounced in obesity (**Fig. 1**). The prevalence of obesity in children of low-income households was 2.3 times greater in boys and 2.2 times greater in girls compared to higher-income households, while the prevalence of overweight was 1.2 times greater in girls of low-income households, without significant differences in boys. In general, for each income bracket, overweight was more frequent than obesity, except in children from low-income households, in whom obesity was more frequent (**Appendix A, Supplemental Table A1**).

The prevalence of central obesity in the lowest-income bracket nearly doubled the prevalence in the highest-income bracket in both boys and girls (**Fig. 2**). The assessment of the BMI and the WtHR as continuous variables also revealed an inverse association between both these variables and income level (**Appendix A, Supplemental Figures A1 and A2**).

In the analysis by parental educational attainment, the prevalence of obesity was nearly double in children of

**Table 1** Sociodemographic and family characteristics, lifestyle habits and school environment of participants of the ALADINO 2019 study by sex.

	Boys (n = 8,513)		Girls (n = 8,152)		P
	n	%	n	%	
<i>Age (years)</i>					
6.0–6.9	2,054	24.1%	2,024	23.8%	.153
7.0–7.9	2,254	26.5%	2,033	23.9%	
8.0–8.9	2,257	26.5%	2,193	25.8%	
9.0–9.9	1,948	22.9%	1,902	22.3%	
<i>Weight status of child (WHO)</i>					
Normal weight	4,795	56.3%	4,782	56.2%	<.001
Overweight	1,924	22.6%	2,046	24%	
Obesity	1,704	20%	1,256	14.8%	
<i>Central obesity</i>					
No	6,611	77.7%	6,261	73.6%	.175
Yes	1,896	22.3%	1,888	22.2%	
<i>Household income (euro/year)</i>					
<12,000	1,093	12.8%	1,062	12.5%	.128
12,000–18,000	1,268	14.9%	1,224	14.4%	
18,001–30,000	1,612	18.9%	1,635	19.2%	
30,001–42,000	1,167	13.7%	1,097	12.9%	
>42,000	1,662	19.5%	1,480	17.4%	
<i>Parental educational attainment (highest)</i>					
Primary education	479	5.6%	522	6.1%	.061
Secondary education	3,898	45.8%	3,771	44.3%	
University education	4,003	47%	3,738	43.9%	
<i>Parental employment status (paid employment)</i>					
None employed	568	6.7%	541	6.4%	.695
One employed	2,545	29.9%	2,485	29.2%	
Both employed	5,323	62.5%	5,048	59.3%	
<i>Parental nationality</i>					
Both Spanish citizens	6,292	73.9%	6,052	71.1%	.771
Both EU citizens	496	5.8%	496	5.8%	
Both non-EU citizens	665	7.8%	664	7.8%	
Mixed	717	8.4%	669	7.9%	
<i>Parental obesity</i>					
Neither parent	6,571	77.2%	6,320	74.2%	.514
One or both parents	1,661	19.5%	1,557	18.3%	
<i>History of breastfeeding</i>					
Never or <1 month	2,403	28.2%	2,345	27.5%	.508
≥1 month	5,893	69.2%	5,621	66%	
<i>Underestimation of child weight status by parents</i>					
No	4,459	52.4%	4,714	55.4%	<.001
Yes	3,853	45.3%	3,288	38.6%	
<i>COSI unhealthy dietary habits questionnaire score</i>					
Low	2,528	29.7%	2,291	26.9%	.106
Medium	4,984	58.5%	4,830	56.7%	
High	314	3.7%	321	3.8%	
<i>Physical activity and sedentary behaviour</i>					
Active and not sedentary	4,365	51.3%	3,540	41.6%	<.001
Active and sedentary	1,387	16.3%	1,268	14.9%	
Inactive and not sedentary	1,287	15.1%	1,831	21.5%	
Inactive and sedentary	426	5%	564	6.6%	
<i>Number of screens in room</i>					
None	5,918	69.5%	5,878	69.1%	<.001
One	1,425	16.7%	1,582	18.6%	
Two or more	907	10.7%	425	5%	

Table 1 (Continued)

	Boys (n = 8,513)		Girls (n = 8,152)		P
	n	%	n	%	
<i>Type of school (ownership)</i>					
Private (including publicly funded)	2,978	35%	2,805	33%	.437
Public	5,535	65%	5,347	62.8%	
<i>Availability of school food services</i>					
Yes	7,706	90.5%	7,395	86.9%	.668
No	807	9.5%	757	8.9%	
<i>Type of school food services</i>					
Onsite kitchen	4,000	47%	3,852	45.3%	.771
External catering	3,596	42.2%	3,430	40.3%	
<i>Availability of extracurricular sports activities</i>					
Yes	6,248	73.4%	5,932	69.7%	.320
No	2,143	25.2%	2,108	24.8%	
<i>Availability of indoor sports facilities</i>					
Yes	6,940	81.5%	6,594	77.5%	.295
No	1,573	18.5%	1,558	18.3%	

EU, European Union; WHO, World Health Organization.





Figure 1 Weight status (WHO standards) of participants in the ALADINO 2019 study by household income and sex of the child.

parents with low educational attainment compared to children of parents with a university education for both sexes (Appendix A, Supplemental Table A1). The trends for central obesity had the same direction but were not as pronounced. In relation to parental employment status, the prevalence of obesity in children of either sex was 1.4 times greater when neither parent was employed compared to when both parents had paid employment. In the case of central obesity, we found no differences in girls, but there were differences in boys (1.3 times increase) in the same direction (Appendix A, Supplemental Table A2).

### Association between family characteristics and household income

For both sexes, there was a direct association between parental educational attainment and household income, and the frequency of a university education was seven-fold in the highest income bracket compared to the lowest (Table 2). In most middle- or high-income households, one or both parents were employed, while in low-income households it was more frequent for a single parent to be employed.



**Figure 2** Central obesity (waist-to-height ratio >0.5) of participants in the ALADINO 2019 study by household income and sex of the child.

On the other hand, in the highest-income bracket, 92% of schoolchildren had Spanish parents, a percentage that nearly doubled the percentage in the lowest-income bracket. The probability of having at least one obese parent was greater in low-income households (2.1 times greater in boys and 1.7 in girls) compared to high-income households. The history of breastfeeding was 1.1 times more frequent in boys and girls from high-income households compared to children from low-income households.

### Association between perceptions and lifestyle habits and household income

Parents of low socioeconomic status were more likely to underestimate the weight status of schoolchildren compared to those of high socioeconomic status (Table 3). We also found an inverse association between the score in the COSI questionnaire of dietary habits and income: unhealthy dietary habits (higher scores) were 6 times more prevalent in schoolchildren from low-income households compared to high-income households.

Physical activity and sedentary behaviour were directly and inversely associated, respectively, to household income level: the most sedentary and least physically active children were more likely to come from lower-income households. In every income bracket, the frequency of physical activity and lack of sedentary behaviour was about 10 points higher in boys compared to girls.

The number of screens in the student's room was inversely correlated to income: having 2 or more screens in the room was nearly 4 times more frequent in boys and 5 times more frequent in girls from low-income households, compared to high-income households. We also ought to highlight that the presence of electronic devices in the room was more common in boys than girls.

### Association between school characteristics and household income

Attendance to private or publicly funded private schools was twice as frequent in students from high-income homes compared to low-income homes (Table 4). Access to school lunch services and attendance to schools where the food was cooked on-site versus brought by a catering service were more frequent among students from high-income compared to low-income households (Table 4). Similarly, attendance to schools that organised sports activities or with indoor sports facilities was more frequent in students of high socioeconomic status.

### Discussion

In our study, we made a detailed analysis of the inverse association between childhood obesity and overweight, on one hand, and socioeconomic status, on the other, considering several mediating factors, thus offering a thorough and representative nationwide analysis of the social and gender gradient of weight status in the paediatric population and associated environmental variables for the first time.

The inverse gradient observed between excess weight (both obesity and overweight) and socioeconomic status, which was more evident for obesity than for overweight and for girls than for boys, was consistent with the findings of other studies conducted in Spain<sup>13</sup> and at the international level.<sup>20</sup> In addition, the data from the ALADINO study show diverging trends based on socioeconomic status: between 2015<sup>11</sup> and 2019,<sup>6</sup> the prevalence of overweight and of obesity increased in low-income households (by 0.9% and 0.5%, respectively) and the prevalence of overweight in middle-income households (by 1.2%), while in households in the highest income bracket, the prevalence of overweight decreased by 1.5% and the prevalence of obesity by 0.8%.



**Table 2** Sociodemographic and family characteristics by annual household income and sex of the child in the ALADINO 2019 study participants.

	Boys % (95% CI)				
	<12,000 €	12,000–18,000 €	18,001–30,000 €	30,001–42,000 €	>42,000 €
<i>Parental educational attainment</i>					
Primary education	15.9 (13.5–18.6)	8.4 (6.8–10.3)	3.8 (2.9–5.1)	0.9 (0.5–1.8)	0.3 (0.1–0.9)
Secondary education	71.2 (67.9–74.2)	68.3 (65.3–71.3)	56.4 (53.5–59.2)	32.5 (29.4–35.8)	11.3 (9.6–13.3)
University education	13 (10.8–15.5)	23.2 (20.6–26.1)	39.8 (37–42.6)	66.6 (63.2–69.7)	88.3 (86.3–90.1)
<i>Parental employment status (paid employment)</i>					
None employed	20.9 (18.2–23.9)	6.1 (4.7–7.9)	1.1 (0.7–1.9)	0.4 (0.2–1)	0.5 (0.2–1.1)
One employed	51.9 (48.4–55.4)	49.8 (46.6–53)	28.3 (25.9–31)	18.1 (15.6–20.8)	12 (10.2–14)
Both employed	27.2 (24.2–30.5)	44.1 (40.9–47.3)	70.5 (67.9–73)	81.5 (78.7–84)	87.5 (85.5–89.3)
<i>Parental nationality</i>					
Both Spanish citizens	48.1 (44.5–51.8)	65.2 (62–68.3)	78.6 (76.1–81)	86.5 (83.9–88.7)	92 (90.3–93.5)
Both EU citizens	16.1 (13.6–19)	10.1 (8.3–12.3)	4.3 (3.3–5.7)	1.5 (0.9–2.6)	0.6 (0.3–1.3)
Both non-EU citizens	24.1 (21.1–27.4)	14.8 (12.6–17.3)	6.5 (5.2–8.2)	3.2 (2.1–4.7)	0.9 (0.5–1.7)
Mixed	11.6 (9.5–14.2)	9.8 (8.1–11.9)	10.5 (8.8–12.5)	8.8 (7–11)	6.4 (5.1–8)
<i>Parental weight status: obesity</i>					
Neither parent	72.4 (69.1–75.4)	74.7 (71.8–77.4)	79.9 (77.5–82.1)	84 (81.4–86.3)	86.3 (84.2–88.1)
One or both parents	27.6 (24.6–30.9)	25.3 (22.6–28.2)	20.1 (17.9–22.5)	16 (13.7–18.6)	13.7 (11.9–15.8)
<i>History of breastfeeding</i>					
Never or <1 month	32.8 (29.5–36.1)	32.7 (29.8–35.8)	31 (28.4–33.7)	27.7 (24.8–30.9)	23.3 (21–25.9)
≥1 month	67.2 (63.9–70.5)	67.3 (64.2–70.2)	69 (66.3–71.6)	72.3 (69.1–75.2)	76.7 (74.1–79)
	Girls % (95 % CI)				
	<12,000 €	12,000–18,000 €	18,001–30,000 €	30,001–42,000 €	>42,000 €
<i>Parental educational attainment</i>					
Primary education	20.8 (18.1–23.9)	9.8 (8.1–11.9)	3.7 (2.7–5)	1.3 (0.7–2.5)	0.4 (0.1–1)
Secondary education	66.7 (63.3–70)	70.2 (67.1–73.1)	55.5 (52.6–58.3)	32.2 (29–35.6)	12.4 (10.6–14.6)
University education	12.4 (10.3–15)	20 (17.5–22.7)	40.8 (38–43.7)	66.4 (63–69.7)	87.2 (85–89.1)
<i>Parental employment status (paid employment)</i>					
None employed	20.8 (18–23.9)	4.1 (3–5.6)	1.4 (0.8–2.2)	1.2 (0.6–2.4)	0.7 (0.3–1.5)
One employed	52 (48.3–55.6)	53 (49.8–56.3)	29.9 (27.4–32.6)	16.6 (14.2–19.3)	11 (9.2–13.1)
Both employed	27.2 (24–30.6)	42.8 (39.7–46.1)	68.7 (66–71.3)	82.2 (79.4–84.6)	88.3 (86.2–90.1)
<i>Parental nationality</i>					
Both Spanish citizens	51.6 (47.9–55.3)	64.8 (61.6–67.9)	79.2 (76.7–81.5)	90.2 (87.9–92.1)	91.5 (89.6–93)
Both EU citizens	14 (11.6–16.7)	10.8 (8.9–13)	4.9 (3.7–6.3)	0.8 (0.4–1.8)	1 (0.6–1.8)
Both non-EU citizens	22.4 (19.4–25.6)	13.9 (11.7–16.4)	6.1 (4.8–7.6)	2.4 (1.5–3.8)	1.2 (0.7–2.1)
Mixed	12.1 (9.9–14.7)	10.5 (8.6–12.8)	9.9 (8.3–11.8)	6.6 (5–8.5)	6.3 (4.9–7.9)
<i>Parental weight status: obesity</i>					
Neither parent	76.9 (73.7–79.8)	72.8 (69.8–75.6)	80.5 (78.1–82.7)	84 (81.3–86.4)	86.2 (83.9–88.2)
One or both parents	23.1 (20.2–26.3)	27.2 (24.4–30.2)	19.5 (17.3–21.9)	16 (13.6–18.7)	13.8 (11.8–16.1)
<i>History of breastfeeding</i>					
Never or <1 month	32.9 (29.6–36.4)	30.1 (27.2–33.2)	33.9 (31.2–36.7)	26.7 (23.7–30)	24.4 (21.9–27.2)
≥1 month	67.1 (63.6–70.4)	69.9 (66.8–72.8)	66.1 (63.3–68.8)	73.3 (70–76.3)	75.6 (72.8–78.1)

CI, confidence interval; EU, European Union.

These differing trends in weight status based on income level have also been observed in previous studies in Spain<sup>21</sup> and other European countries.<sup>22</sup>

Our findings show that the association between poverty and obesity is not only present in schoolchildren, but also in their parents. A previous study conducted in Spain also found an increased risk of obesity in children whose parents (one or both) had obesity, and this intergenerational pattern was more pronounced in boys than in girls and in the households in the extremes of the income distri-

bution by quartiles.<sup>21</sup> We also found a direct association between income and the history of breastfeeding.<sup>23</sup> In families of low socioeconomic status, this may be due to a lower maternal educational attainment and poorer working conditions.<sup>24,25</sup>

Another relevant finding was that parents perceived the weight status of the child incorrectly more frequently in low-income households. Once more, these findings could be explained by the lower educational attainment of these parents and/or an increased prevalence of obesity among them,

**Table 3** Parental perceptions and schoolchildren habits by income level and child sex in the ALADINO 2019 sample.

	Boys % (95% CI)				
	<12,000 €	12,000–18,000 €	18,001–30,000 €	30,001–42,000 €	>42,000 €
<i>Parental underestimation of the weight status of the child</i>					
No	53 (49.4–56.4)	48.4 (45.2–51.7)	53.1 (50.3–56)	56 (52.6–59.4)	63.4 (60.6–66.1)
Yes	47 (43.6–50.6)	51.6 (48.3–54.8)	46.9 (44–49.7)	44 (40.6–47.4)	36.6 (33.9–39.4)
<i>COSI unhealthy dietary habits score<sup>a</sup></i>					
Low	17.2 (14.6–20.2)	22.6 (19.9–25.5)	28.9 (26.3–31.6)	37.3 (34–40.8)	45.3 (42.4–48.3)
Medium	74.3 (70.9–77.4)	72.3 (69.2–75.2)	66.3 (63.5–69)	61.2 (57.7–64.5)	53.5 (50.6–56.4)
High	8.5 (6.6–10.8)	5.1 (3.8–6.8)	4.8 (3.7–6.2)	1.5 (0.8–2.6)	1.2 (0.7–2)
<i>Physical activity and sedentary behaviour</i>					
Active and not sedentary	46.1 (42.4–49.8)	51.4 (48–54.8)	56.9 (53.9–59.9)	63.4 (59.9–66.8)	64.4 (61.5–67.3)
Active and sedentary	26.5 (23.3–29.9)	24.1 (21.3–27.2)	20.2 (17.9–22.7)	14.5 (12.2–17.2)	9.6 (7.9–11.5)
Inactive and not sedentary	19.5 (16.7–22.8)	15 (12.7–17.6)	17.2 (15–19.6)	18.6 (15.9–21.6)	22.3 (19.8–24.9)
Inactive and sedentary	7.9 (6.1–10.2)	9.4 (7.6–11.7)	5.7 (4.5–7.2)	3.5 (2.4–5)	3.7 (2.7–5.1)
<i>Number of screens in room</i>					
None	56.1 (52.6–59.6)	61.2 (57.9–64.3)	68.9 (66.1–71.5)	78.7 (75.7–81.3)	85.6 (83.4–87.6)
One	25.1 (22.2–28.3)	21.8 (19.2–24.6)	18.1 (16–20.5)	13.5 (11.3–16)	9.2 (7.7–11.1)
Two or more	18.7 (16.1–21.7)	17.1 (14.7–19.7)	13 (11.1–15.1)	7.8 (6.2–9.9)	5.1 (4–6.6)
	Girls % (95% CI)				
	<12,000 €	12,000–18,000 €	18,001–30,000 €	30,001–42,000 €	>42,000 €
<i>Parental underestimation of the weight status of the child</i>					
No	50.9 (47.3–54.5)	56.2 (53–59.4)	57.5 (54.6–60.3)	66.2 (62.8–69.5)	67.1 (64.2–69.9)
Yes	49.1 (45.5–52.7)	43.8 (40.6–47)	42.5 (39.7–45.4)	33.8 (30.5–37.2)	32.9 (30.1–35.8)
<i>COSI unhealthy dietary habits score<sup>a</sup></i>					
Low	18.7 (15.8–21.9)	23.2 (20.4–26.2)	25.4 (23–28.1)	35.4 (32–38.9)	46.9 (43.8–50.1)
Medium	70.9 (67.2–74.3)	71.1 (67.9–74.1)	69.8 (67–72.4)	62.1 (58.5–65.5)	51.3 (48.2–54.4)
High	10.4 (8.2–13)	5.7 (4.3–7.5)	4.8 (3.7–6.3)	2.5 (1.6–4)	1.7 (1.1–2.8)
<i>Physical activity and sedentary behaviour</i>					
Active and not sedentary	38.3 (34.6–42)	43.1 (39.7–46.5)	47.8 (44.8–50.8)	53.4 (49.7–57.1)	53.5 (50.3–56.7)
Active and sedentary	26.6 (23.3–30.1)	25.5 (22.6–28.6)	18.7 (16.5–21.3)	12.9 (10.6–15.6)	7.9 (6.3–9.8)
Inactive and not sedentary	25.3 (22.1–28.9)	20.6 (18–23.5)	24.5 (21.9–27.2)	26.7 (23.5–30.1)	34.1 (31.1–37.2)
Inactive and sedentary	9.8 (7.8–12.3)	10.8 (8.8–13.3)	9 (7.4–10.9)	7.1 (5.4–9.2)	4.5 (3.4–6.1)
<i>Number of screens in room</i>					
None	65 (61.4–68.5)	61.5 (58.3–64.7)	71.8 (69.2–74.4)	82.5 (79.6–85.1)	87.1 (84.9–89)
One	26.2 (23.1–29.6)	30.6 (27.7–33.8)	22.6 (20.3–25.1)	13.5 (11.2–16.1)	11.1 (9.3–13.2)
Two or more	8.8 (6.9–11.1)	7.8 (6.2–9.8)	5.5 (4.4–7)	4 (2.8–5.7)	1.8 (1.2–2.9)

CI, confidence interval; COSI, Childhood Obesity Surveillance Initiative.

<sup>a</sup> COSI unhealthy dietary habits score: one point given for each of the contemplated unhealthy dietary habits (skipping breakfast, failure to consume fruit daily, failure to consume vegetables daily, consumption of sugary drinks 4 or more times a week, consumption of sweet snacks four or more times a week, consumption of salty snacks four or more times a week). The possible total score ranges from 0 to 6, where 0 is healthiest and 6 least healthy. We categorised the score as low (0–1 points), medium (2–3 points) and high (4–6 points).

which would make them less likely to perceive excess weight in their children as a problem.<sup>26</sup>

The direct association between low household income and unhealthier dietary habits, in agreement with other studies conducted in Spain<sup>27</sup> and at the international level,<sup>28</sup> may be mediated by various mechanisms, such as poorer nutrition literacy in relation to a lower educational attainment, financial difficulties that limit shopping choice (especially of fresh foods) or lacking the time to cook, which in turn are associated with an increased consumption of precooked or processed foods, which have poorer nutrient profiles.<sup>29,30</sup>

Physical activity and sedentary behaviour were also associated with socioeconomic status in our study. We also ought to highlight the observed gender gap: girls were less physically active than boys, even within the same income bracket. The causes of this double inequality are multiple, and chief among them are, on one hand, greater access to sports activities and facilities in higher-income areas<sup>31</sup> and a greater awareness of exercise as a contributor to health in families with greater educational attainment<sup>32</sup> and, on the other, a family environment that is more supportive of sports in households with male offspring.<sup>33</sup>



**Table 4** Characteristics of the school environment by income level and child sex in the ALADINO 2019 sample.

	Boys % (95% CI)				
	<12,000 €	12,000–18,000 €	18,001–30,000 €	30,001–42,000 €	>42,000 €
<i>Type of school (ownership)</i>					
Private (including publicly funded)	19.5 (16.9–22.4)	25.9 (23.2–28.9)	29.9 (27.3–32.5)	35.4 (32.3–38.7)	49.1 (46.2–52)
Public	80.5 (77.6–83.1)	74.1 (71.1–76.8)	70.1 (67.5–72.7)	64.6 (61.3–67.7)	50.9 (48–53.8)
<i>Availability of school food services</i>					
Yes	89.3 (87.1–91.1)	92.2 (90.5–93.6)	92.1 (90.5–93.4)	95 (93.4–96.3)	97.4 (96.4–98.2)
No	10.7 (8.9–12.9)	7.8 (6.4–9.5)	7.9 (6.6–9.5)	5 (3.7–6.6)	2.6 (1.8–3.6)
<i>Type of school food services</i>					
Onsite kitchen	43 (39.4–46.8)	44.4 (41–47.8)	51.9 (48.9–54.9)	55.1 (51.5–58.5)	61.5 (58.6–64.3)
External catering	57 (53.2–60.6)	55.6 (52.2–59)	48.1 (45.1–51.1)	44.9 (41.5–48.5)	38.5 (35.7–41.4)
<i>Availability of extracurricular sports activities</i>					
Yes	75.1 (72–77.9)	74.4 (71.6–77.1)	73.5 (70.9–75.9)	80.3 (77.5–82.8)	82.6 (80.3–84.6)
No	24.9 (22.1–28)	25.6 (22.9–28.4)	26.5 (24.1–29.1)	19.7 (17.2–22.5)	17.4 (15.4–19.7)
<i>Availability of indoor sports facilities</i>					
Yes	76.5 (73.5–79.3)	77.2 (74.4–79.7)	78.4 (76–80.7)	78.9 (76–81.6)	83 (80.7–85.1)
No	23.5 (20.7–26.5)	22.8 (20.3–25.6)	21.6 (19.3–24)	21.1 (18.4–24)	17 (14.9–19.3)
	Girls % (95% CI)				
	<12,000 €	12,000–18,000 €	18,001–30,000 €	30,001–42,000 €	>42,000 €
<i>Type of school (ownership)</i>					
Private (including publicly funded)	21.4 (18.6–24.5)	23.2 (20.6–26.1)	30.8 (28.2–33.5)	34.2 (31–37.6)	47.8 (44.7–50.8)
Public	78.6 (75.5–81.4)	76.8 (73.9–79.4)	69.2 (66.5–71.8)	65.8 (62.4–69)	52.2 (49.2–55.3)
<i>Availability of school food services</i>					
Yes	92.3 (90.4–93.8)	89.2 (87.2–91)	93.6 (92.1–94.8)	94.3 (92.5–95.7)	97 (95.9–97.8)
No	7.7 (6.2–9.6)	10.8 (9–12.8)	6.4 (5.2–7.9)	5.7 (4.3–7.5)	3 (2.2–4.1)
<i>Type of school food services</i>					
Onsite kitchen	45.1 (41.4–48.9)	46.1 (42.6–49.6)	52.7 (49.7–55.7)	55.1 (51.4–58.7)	60.7 (57.6–63.7)
External catering	54.9 (51.1–58.6)	53.9 (50.4–57.4)	47.3 (44.3–50.3)	44.9 (41.3–48.6)	39.3 (36.3–42.4)
<i>Availability of extracurricular sports activities</i>					
Yes	73.2 (70–76.2)	76.3 (73.5–78.9)	74.6 (72.1–77)	79 (76–81.7)	81.6 (79.1–83.8)
No	26.8 (23.8–30)	23.7 (21.1–26.5)	25.4 (23–27.9)	21 (18.3–24)	18.4 (16.2–20.9)
<i>Availability of indoor sports facilities</i>					
Yes	75.5 (72.4–78.4)	77.3 (74.5–79.9)	76.7 (74.1–79)	79.2 (76.2–81.9)	83.3 (80.9–85.5)
No	24.5 (21.6–27.6)	22.7 (20.1–25.5)	23.3 (21–25.9)	20.8 (18.1–23.8)	16.7 (14.5–19.1)

CI, confidence interval.

The greater presence of electronic devices in the rooms of schoolchildren in households of lower socioeconomic status has also been observed in neighbouring countries.<sup>34</sup> Parental educational attainment, parental setting of screentime limits and parental screentime could explain the observed differences based on income level.<sup>35</sup>

Other factors that were not addressed in our study are also associated with household socioeconomic status and the risk of childhood obesity, such as family structure, understood as the number of members of the household and the relationship between them, with evidence of an increased risk of obesity in households with a single parent and house-

holds with a single child.<sup>36</sup> Another aspect that has been explored in the literature that is also associated with socioeconomic status and whose prevalence is increasing in the paediatric population is the presence of mental health problems in the child.<sup>37</sup>

When it came to the school environment, our study found that income level was also related to the type of school attended by the child as well as the available facilities and services. Indeed, the availability and actual access to sports facilities in the school are among the factors that may contribute to the level of physical activity in school-aged children, since, for instance, the number of facilities devoted to sports and play in the school has been found to

be directly associated with the level of light physical activity and inversely associated with sedentary behaviour.<sup>38</sup>

Based on the results of the study, schoolchildren from low-income households grow in household and school environments that increase the risk of obesity, giving rise to what is known as an obesogenic environment through the interaction of factors of a very diverse nature: parental excess weight, lack of breastfeeding, inaccurate perception of the child's weight status by the parents, less healthy lifestyle habits in terms of nutrition, physical activity and sedentary behaviour, greater availability of electronic devices and a school environment lacking a kitchen of its own, extracurricular sports activities or indoor sports facilities. Our findings highlight the need to develop effective multicomponent interventions to decrease childhood obesity, a problem exacerbated by the double income and gender gap, on the different environments.

The current literature supports the effectiveness of some early interventions to address both childhood obesity and child poverty.<sup>39</sup> In Spain, a recent study found a lower prevalence of childhood obesity in regions where more resources were devoted to public health interventions, although prevention policy tended to have a greater effect in middle- and high-income groups, thus reproducing existing health inequalities.<sup>40</sup>

To address the inequalities and barriers identified in the study, it would be advisable to implement strategies aimed at reducing the burden of the obesogenic environment, and schools can offer an optimal equalising mechanism to narrow the income and gender gaps. The most important strategies would be those aimed at promoting a healthy diet and physical activity and those aimed at strengthening parental knowledge and awareness, especially in the most disadvantaged environments, in addition to ongoing epidemiological surveillance of childhood obesity and its determinants, with a particular emphasis on inequities.

The main strength of the study was that the analysis with a sex-gender perspective of the data of the ALADINO study, conducted in a large sample with objective anthropometric measurements and collection of data on multiple factor associated with the schoolchildren, their family background and their school environment, allowing the study of the obesogenic environment as a whole. Among its limitations, we ought to highlight that the data collected through self-report questionnaires may be a source of information biases such as recall bias and that the cross-sectional design of the study precluded the establishment of causality in the identified associations.

## Conclusion

The prevalences of both overweight and obesity in schoolchildren in Spain are inversely associated to household socioeconomic status. Several factors, such as dietary and physical activity habits, the family background and the school environment, are also strongly associated with family socioeconomic status and may act as mediators in the aforementioned association, and there are also relevant differences based on the sex of the child. Interventions aimed at this important public health problem must be prioritised and adapted taking into account the identified inequalities

to ensure their implementation can reach all children and be equally effective regardless of their socioeconomic background.

## Funding

The ALADINO 2019 study was funded by the Agencia Española de Seguridad Alimentaria y Nutrición, which carries rounds of the ALADINO study at regular intervals in the framework of the Childhood Obesity Surveillance Initiative of the WHO Regional Office for Europe.

## Conflicts of interest

The authors have no conflicts of interest to declare.

## Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.anpedi.2023.05.013>.

## References

1. Kumar S, Kelly AS. Review of childhood obesity: from epidemiology, etiology, and comorbidities to clinical assessment and treatment. *Mayo Clin Proc.* 2017;92:251–65.
2. Herouvi D, Karanasios E, Karayianni C, Karavanaki K. Cardiovascular disease in childhood: the role of obesity. *Eur J Pediatr.* 2013;172:721–32.
3. Simmonds M, Llewellyn A, Owen CG, Woolacott N. Predicting adult obesity from childhood obesity: a systematic review and meta-analysis. *Obes Rev Off J Int Assoc Study Obes.* 2016;17:95–107.
4. Nittari G, Scuri S, Petrelli F, Pirillo I, di Luca NM, Grappasonni I. Fighting obesity in children from European World Health Organization member states. *Epidemiological data, medical-social aspects, and prevention programs.* *Clin Ter.* 2019;170:e223–30.
5. Report on the fifth round of data collection, 2018–2020: WHO European Childhood Obesity Surveillance Initiative (COSI) [Internet]. [citado 15 de noviembre de 2022]. Available from: <https://www.who.int/europe/publications/i/item/WHO-EURO-2022-6594-46360-67071>.
6. García-Solano M, Gutiérrez-González E, López-Sobaler AM, Ruiz-Álvarez M, Bermejo López LM, Aparicio A, et al. Situación ponderal de la población escolar de 6 a 9 años en España: resultados del estudio ALADINO 2019. *Nutr Hosp.* 2021;38:943–53.
7. WHO Regional Office for Europe; WHO European Regional Obesity Report 2022 [Internet]. [citado 22 de diciembre de 2022]. Available from: <https://apps.who.int/iris/bitstream/handle/10665/353747/9789289057738-eng.pdf>.
8. Chaudry A, Wimer C. Poverty is not just an indicator: the relationship between income, poverty, and child well-being. *Acad Pediatr.* 2016;16:S23–29.
9. Gobierno España - Alto Comisionado para la lucha contra la pobreza infantil. Tasas de riesgo de pobreza infantil en España (2008-2020) [Internet]. 2021 [citado 22 de diciembre de 2022]. Available from: <https://www.comisionadopobrezainfantil.gob.es/es/tasas-de-riesgo-de-pobreza-infantil-en-espa%C3%B1a-2008-2020>.
10. Brown T, Moore TH, Hooper L, Gao Y, Zayegh A, Ijaz S, et al. Interventions for preventing obesity in children. *Cochrane Database Syst Rev.* 2019;7:CD001871.

11. García-Solano M, Gutiérrez-González E, López-Sobaler AM, Dal Re Saavedra MÁ, Robledo de Dios T, Villar-Villalba C, et al. Weight status in the 6 to 9 year-old school population in Spain: results of the ALADINO 2015 study. *An Pediatr.* 2021;94:366–76.
12. Pérez-Farinós N, López-Sobaler AM, Dal Re MÁ, Villar C, Labrado E, Robledo T, et al. The ALADINO study: a national study of prevalence of overweight and obesity in Spanish children in 2011. *BioMed Res Int.* 2013;2013:163687.
13. Ministerio de Sanidad, Consumo y Bienestar Social - Portal Estadístico del SNS - Encuesta Nacional de Salud de España 2017 [Internet]. [citado 22 de diciembre de 2022]. Available from: <https://www.msbs.gob.es/estadEstudios/estadisticas/encuestaNacional/encuesta2017.htm>.
14. Childhood Obesity Surveillance Initiative (COSI) protocol (2016) [Internet]. [citado 22 de diciembre de 2022]. Available from: <https://apps.who.int/iris/handle/10665/354793>.
15. Childhood Obesity Surveillance Initiative (COSI) - Data collection procedures (2016) [Internet]. [citado 22 de diciembre de 2022]. Available from: <https://apps.who.int/iris/handle/10665/354792>.
16. Agencia Española de Seguridad Alimentaria y Nutrición. Estudio ALADINO 2019: Estudio sobre la Alimentación, Actividad Física, Desarrollo Infantil y Obesidad en España 2019 [Internet]. 2019. Available from: [https://www.aesan.gob.es/AECOSAN/docs/documentos/nutricion/observatorio/Informe\\_Aladino\\_2019.pdf](https://www.aesan.gob.es/AECOSAN/docs/documentos/nutricion/observatorio/Informe_Aladino_2019.pdf).
17. WHO | Growth reference data for 5-19 years [Internet]. WHO. [citado 22 de diciembre de 2022]. Available from: <https://www.who.int/tools/growth-reference-data-for-5to19-years>.
18. Taxová Braunerová R, Kunešová M, Heinen MM, Rutter H, Hassapidou M, Duleva V, et al. Waist circumference and waist-to-height ratio in 7-year-old children-WHO Childhood Obesity Surveillance Initiative. *Obes Rev Off J Int Assoc Study Obes.* 2021;22 Suppl 6:e13208.
19. Williams J, Buoncristiano M, Nardone P, Rito AI, Spinelli A, Hejgaard T, et al. A snapshot of European children's eating habits: results from the fourth round of the WHO European Childhood Obesity Surveillance Initiative (COSI). *Nutrients.* 2020;12:E2481.
20. Buoncristiano M, Williams J, Simmonds P, Nurk E, Ahrens W, Nardone P, et al. Socioeconomic inequalities in overweight and obesity among 6- to 9-year-old children in 24 countries from the World Health Organization European region. *Obes Rev Off J Int Assoc Study Obes.* 2021;22 Suppl 6:e13213.
21. Costa-Font J, Gil J. Intergenerational and socioeconomic gradients of child obesity. *Soc Sci Med* 1982. 2013;93:29–37.
22. Bouthoorn SH, Wijtzes AI, Jaddoe VWV, Hofman A, Raat H, van Lenthe FJ. Development of socioeconomic inequalities in obesity among Dutch pre-school and school-aged children. *Obes Silver Spring Md.* 2014;22:2230–7.
23. Uwaezuoke SN, Eneh CI, Ndu IK. Relationship between exclusive breastfeeding and lower risk of childhood obesity: a narrative review of published evidence. *Clin Med Insights Pediatr.* 2017;11:1179556517690196.
24. Villar M, Santa-Marina L, Murcia M, Amiano P, Gimeno S, Ballester F, et al. Social factors associated with non-initiation and cessation of predominant breastfeeding in a mother-child cohort in Spain. *Matern Child Health J.* 2018;22:725–34.
25. Gutierrez-de-Terán-Moreno G, Ruiz-Litago F, Ariz U, Fernández-Atutxa A, Mulas-Martín MJ, Benito-Fernández E, et al. Successful breastfeeding among women with intention to breastfeed: From physiology to socio-cultural factors. *Early Hum Dev.* 2022;164:105518.
26. Blanchet R, Kengneson CC, Bodnaruc AM, Gunter A, Giroux I. Factors influencing parents' and children's misperception of children's weight status: a systematic review of current research. *Curr Obes Rep.* 2019;8:373–412.
27. Miqueleiz E, Lostao L, Ortega P, Santos JM, Astasio P, Regidor E. Patrón socioeconómico en la alimentación no saludable en niños y adolescentes en España. *Aten Primaria.* 2014;46:433–9.
28. Fismen AS, Buoncristiano M, Williams J, Helleve A, Abdrahmanova S, Bakacs M, et al. Socioeconomic differences in food habits among 6- to 9-year-old children from 23 countries-WHO European Childhood Obesity Surveillance Initiative (COSI 2015/2017). *Obes Rev Off J Int Assoc Study Obes.* 2021;22 Suppl 6:e13211.
29. Khanom A, Hill RA, Morgan K, Rapport FL, Lyons RA, Brophy S. Parental recommendations for population level interventions to support infant and family dietary choices: a qualitative study from the Growing Up in Wales, Environments for Healthy Living (EHL) study. *BMC Public Health.* 2015;15:234.
30. Lindsay AC, Wallington SF, Lees FD, Greaney ML. Exploring how the home environment influences eating and physical activity habits of low-income, latino children of predominantly immigrant families: a qualitative study. *Int J Environ Res Public Health.* 2018;15:978.
31. Anselma M, Chinapaw MJM, Altenburg TM. Determinants of child health behaviors in a disadvantaged area from a community perspective: a participatory needs assessment. *Int J Environ Res Public Health.* 2018;15:644.
32. Seabra A, Mendonça D, Maia J, Welk G, Brustad R, Fonseca AM, et al. Gender, weight status and socioeconomic differences in psychosocial correlates of physical activity in schoolchildren. *J Sci Med Sport.* 2013;16:320–6.
33. Revuelta LR, Etxaniz IE. Clima familiar deportivo y auto-concepto físico en la adolescencia. *Eur J Educ Psychol.* 2011;4:19–31.
34. Rodrigues D, Gama A, Machado-Rodrigues AM, Nogueira H, Rosado-Marques V, Silva MRG, et al. Home vs. bedroom media devices: socioeconomic disparities and association with childhood screen- and sleep-time. *Sleep Med.* 2021;83:230–4.
35. Salway RE, Emm-Collison L, Sebire S, Thompson JL, Jago R. Associations between socioeconomic position and changes in children's screen-viewing between ages 6 and 9: a longitudinal study. *BMJ Open.* 2019;9:e027481.
36. Kanellopoulou A, Giannakopoulou SP, Notara V, Antonogeorgos G, Rojas-Gil AP, Kornilaki EN, et al. The association between adherence to the Mediterranean diet and childhood obesity; the role of family structure: results from an epidemiological study in 1728 Greek students. *Nutr Health.* 2021;27:39–47.
37. Förster LJ, Vogel M, Stein R, Hilbert A, Breinker JL, Böttcher M, et al. Mental health in children and adolescents with overweight or obesity. *BMC Public Health.* 2023;23:135.
38. Dalene KE, Anderssen SA, Ekelund U, Thorén AKH, Hansen BH, Kolle E. Permanent play facility provision is associated with children's time spent sedentary and in light physical activity during school hours: a cross-sectional study. *Prev Med Rep.* 2016;4:429–34.
39. Taghizadeh S, Farhangi MA. The effectiveness of pediatric obesity prevention policies: a comprehensive systematic review and dose-response meta-analysis of controlled clinical trials. *J Transl Med.* 2020;18:480.
40. Carmona-Rosado L, Zapata-Moya ÁR. Los esfuerzos preventivos de las comunidades autónomas y la desigualdad socioeconómica en la obesidad o el sobrepeso infantil. *Gac Sanit.* 2022;36:214–20.